

Piecewise Stereoselective Assembly of Multisubstituted Alkenes

Eli Jones,[‡] Robert T. Martin,[‡] and David W. C. MacMillan*Cite This: *J. Am. Chem. Soc.* 2026, 148, 23499–23505

Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: Multisubstituted alkenes are valuable structural motifs that also serve as key intermediates in complexity-building transformations; however, their stereoselective synthesis remains a longstanding challenge. Alkenyl bromides are appealing coupling handles for alkene synthesis, though their application would be greatly enhanced if they could be coupled directly with alkyl radicals derived from feedstock alcohols. Herein, we report a nickel metallaphotoredox-enabled deoxygenative cross-coupling of alkenyl bromides with alcohols, providing a direct, modular approach to multisubstituted alkenes. This transformation exhibits a broad substrate scope across both alcohol and alkenyl bromide partners, delivering over 30 multisubstituted alkenes. Moreover, we developed a complementary stereoselective bromoalkenylation from readily synthesized *gem*-dibromoolefins to address the limited availability of stereodefined alkenyl bromides. In the presence of a phthalimide additive, selective monofunctionalization enables access to over 40 stereodefined alkenyl bromides, providing a general and stereoselective pathway to these otherwise difficult-to-access intermediates. Direct access to alkenyl bromides enables piecewise stereoselective assembly of multisubstituted alkenes through sequential alcohol coupling or stereoretentive Suzuki–Miyaura reactions. These strategies are illustrated via iterative cross-coupling sequences and the total synthesis of (+)-sponalisolide B.

Multisubstituted alkenes are ubiquitous motifs in pharmaceuticals, natural products,¹ and functional materials and also serve as versatile synthetic building blocks.^{2,3} Although classical alkene syntheses, including carbonyl olefination⁴ and olefin metathesis,⁵ have been reliably used to access disubstituted alkenes, the stereoselective synthesis of tri- and tetrasubstituted alkenes remains challenging.⁶ Established approaches to these motifs include alkyne difunctionalization,^{7,8} carbonyl olefination,⁹ and elimination-based strategies.¹⁰ However, these methods often fail to offer broad generality across substitution patterns. Cross-coupling has emerged as a powerful strategy for the modular construction of such multisubstituted alkene frameworks. In particular, highly substituted alkenes can be accessed stereoselectively via the direct coupling of structurally diverse vinyl and alkyl or aryl coupling partners.^{11,12} Representative examples include our laboratory's decarboxylative vinylation,^{13,14} Weix's cross-electrophile coupling of vinyl and alkyl halides,¹⁵ and Watson's coupling of tetrasubstituted vinyl silanes with aryl halides.¹⁶ Despite these advances, preparation of the corresponding coupling partners remains challenging: alkyl substrates can be limited in generality while vinyl partners are often difficult to prepare with stereoselectivity.¹²

Recently, our group identified benzoxazolium salt reagents (termed "NHC") as mild, robust activators of alcohols in deoxygenative metallaphotoredox cross-coupling reactions.^{17–22} We reasoned that we could pair this activation strategy with alkenyl bromide electrophiles to achieve alkene synthesis via direct cross-coupling. Given the commercial abundance and structural diversity of alcohols,²³ this approach could enable a modular strategy for the assembly of multisubstituted alkenes from alkenyl bromides (Figure 1B). Alkenyl bromides are typically obtained through multistep

routes from enolizable carbonyl compounds,²⁴ alkynes,²⁵ or functionalized alkenyl fragments.^{26,27} These strategies are often ineffective for all-alkyl alkenyl bromides, where both regio- and stereoselective synthesis can be challenging. To accomplish our desired modular alkene assembly strategy, we also sought a general and stereoselective route to the alkenyl bromide partners themselves.

gem-Dibromoolefins are readily accessible in a single step²⁸ from commercially abundant aldehydes and ketones (Figure 1B). We envisioned that these *gem*-dibromoolefins could serve as suitable electrophiles for selective cross-coupling with alcohol-derived alkyl radicals to furnish alkenyl bromide products, which can serve as intermediates for subsequent, stereocontrolled coupling steps (Figure 1C).^{29–32} If successful, this strategy would provide one of the first general methods to stereoselectively access all-alkyl trisubstituted alkenyl bromides. We recognized two major challenges associated with such a transformation. (1) The reaction must proceed with high chemoselectivity to furnish the alkenyl bromide product, which can itself serve as a competent electrophile. Indeed, Suzuki–Miyaura functionalization of *gem*-dibromoolefins has typically shown limited chemoselectivity, with predominant double coupling observed.^{30–32} (2) The reaction must be stereoselective, functionalizing only one of the two bromides on the substrate.

Received: May 4, 2026

Revised: May 28, 2026

Accepted: June 4, 2026

Published: June 8, 2026






Figure 1. Strategic approach to stereoselective multisubstituted alkene synthesis.

Herein, we overcome both of these challenges using nickel metallaphotoredox catalysis. We further demonstrate that the resulting alkenyl bromide products—and alkenyl bromides more generally—undergo efficient cross-coupling with alkyl radicals or alkylboranes, enabling the piecewise stereoselective assembly of multisubstituted alkenes from aldehydes or ketones and alcohols.

We first set out to develop a general alcohol–alkenyl bromide cross-coupling. Following an initial optimization campaign (see Supporting Information, Figures S01–S03), we identified conditions to couple *tert*-butyl 3-(bromomethylene)azetidine-1-carboxylate with benzyl 4-(2-hydroxyethyl)piperidine-1-carboxylate in 87% yield (Table 1, 20). Encouraged by these results, we evaluated the scope of the transformation with a range of alcohol coupling partners (Table 1). Primary alcohols bearing heterocyclic side chains are well tolerated (4, 5), as are substrates containing alkyl and aryl bromides (6, 1), highlighting selective functionalization of the alkenyl bromide over other potential electrophiles. Allylic and benzylic alcohols (3, 9, 10) are also competent substrates, furnishing skipped diene products that are often challenging to access.³³ The reaction accommodates secondary alcohols, with broad functional-group tolerance for hydridic C–H bonds (11), free alcohols (17), and ketones (18). Complex alcohols are excellent substrates, including sugars (8), steroids (18), and a derivative of the antiplatelet agent ticagrelor (19), all of which undergo vinylation in >70% yield.

We then explored the scope of the alkenyl bromide coupling partner. Alkenyl bromides embedded within six- (21) and


seven-membered rings (22), as well as trisubstituted alkenyl bromides (23, 25), furnish products in good yields. The reaction tolerates pendant functionalities on the alkenyl bromide, including a free alcohol (26) and a trimethylsilyl group (27). We were pleased to find that the deoxygenative vinylation proceeds with retention of stereochemistry for unsymmetric alkenyl bromides, although partial erosion of stereochemical fidelity was observed with (*Z*)-1-bromoprop-1-ene (29). Finally, a 1-bromo-1-fluoroalkene furnishes the fluoroalkene product in 81% yield, with full chemoselectivity for the bromide (31).

Next, we set out to achieve the stereoselective synthesis of alkenyl bromides from *gem*-dibromoolefin precursors, themselves readily prepared from commercially available aldehydes and ketones (see the Supporting Information). Under optimized alcohol–alkenyl bromide coupling conditions (*vide supra*) with increased nickel loading, the alkenyl bromide product was obtained, along with significant amounts of the corresponding dialkylated product (Figure S04). To suppress this undesired pathway, we evaluated additives that could impact the oxidative addition step. Our group has previously shown that phthalimide is an effective additive in nickel-catalyzed metallaphotoredox reactions, where it is proposed to stabilize the oxidative addition adduct and suppress off-cycle reactivity.³⁴ Gratifyingly, we observed that addition of 1.0 equiv of phthalimide both enhanced the yield of the desired alkenyl bromide and suppressed dialkylation (see Supporting Information for further details). We next sought to optimize for product stereoselectivity. Following extensive evaluation of conditions using an aldehyde-derived unsymmetric *gem*-dibromoolefin (51, Figures S05–S10), we found that changing the solvent system to 1:2 MTBE:1,2-dichlorobenzene led to marked improvements in both yield and stereoselectivity.

With optimized conditions in hand, we explored the scope of this transformation (Table 2). Primary alcohols are particularly well-tolerated, including those bearing potentially challenging motifs, such as phosphonate (36), pyridine (38), and trifluoromethyl groups (41). Excitingly, an alcohol substrate bearing an aryl bromide reacted with high efficiency—even in the presence of three potential sites for oxidative addition, only a single stereodefined product is observed (39). Benzylic alcohols are also efficient coupling partners, affording 2-bromoallylarene products (40, 41). A variety of secondary alcohols bearing four- (42), five- (44, 46), six- (43), and seven-membered rings (45) furnish the desired products in good yields. Finally, even complex alcohols, including sugars (47, 50) and alcohol-containing drugs (48, 49), serve as effective alkyl radical sources.

We next examined the scope of the *gem*-dibromoolefin component. Substrates containing tertiary amides (51), thioethers (53), and furans (54) underwent coupling with 4-(2-hydroxyethyl)piperidine-1-carboxylate in good yields, as did conjugated *gem*-dibromoolefins, which efficiently formed the corresponding bromostyrenes (59, 60) and bromoenynes (55). Tertiary substitution at the olefin α -carbon (56, 57) did not meaningfully impair reactivity. Notably, substrates bearing a free carbonyl group were compatible with both the *gem*-dibromoolefin synthesis and subsequent cross-coupling conditions (57). Finally, *gem*-dibromoolefins bearing six- (61), three- (62), and five-membered rings (64) were well-tolerated.

We then applied these conditions to the synthesis of tetrasubstituted alkenes from ketone-derived *gem*-dibromoolefins. A variety of *gem*-dibromoolefins underwent cross-coupling in


Table 1. Alkylation of Alkenyl Bromides^a

^aReactions performed with alcohol (1.75 equiv), alkenyl bromide (0.5 mmol, 1.0 equiv), 5,7-di-*tert*-butyl-3-phenylbenzo[d]oxazol-3-ium tetrafluoroborate (NHC, 1.9 equiv), pyridine (1.9 equiv), quinuclidine (1.9 equiv), [Ir(ppy)₂(dtbbpy)]PF₆ (1.5 mol %), Ni(dtbbpy)Br₂ (5 mol %), MTBE/Acetone (1:1, 0.05 M), integrated photoreactor (450 nm, m1 plate, 100% light intensity), 2 h. ^bAssay yield (by NMR vs. mesitylene as internal standard). ^c7.5 mol % Ni(dtbbpy)Br₂, 0.225 equiv. phthalimide used. ^dd.r. = 1.9:1. ^ed.r. = 1.4:1. ^fd.r. = 5:1. ^gd.r. > 20:1. ^hAlcohol condensation performed in 1,4-dioxane, 1.0 equiv. phthalimide used. ⁱd.r. = 3.2:1. ^j0.225 equiv. phthalimide used. ^k1.0 equiv. phthalimide used. See Supporting Information for experimental details.

good yields, although asymmetric tetra-substituted *gem*-dibromoolefins (**66**, **69**, **70**) suffered erosion of stereoselectivity compared to aldehyde-derived *gem*-dibromoolefins. This may arise from unselective oxidative addition owing to the similar steric environments of the two Br atoms in the *gem*-dibromoolefin, or from an XAT-type oxidative addition³⁵ at more hindered C–Br bonds, producing a vinyl radical that can rapidly isomerize before nickel radical capture.

To showcase the synthetic utility of this transformation, we undertook the piecewise, stereoselective assembly of multi-

substituted alkenes (Figure 2). Stereoselective synthesis of the alkenyl bromide could be followed by a stereoretentive Suzuki–Miyaura cross-coupling¹¹ with an alkylborane to yield *all*-alkyl trisubstituted alkenes. Delightfully, either of the two stereoisomers can be accessed with excellent selectivity by exchanging the alcohol and alkylborane partners ((*E*)-**71**, (*Z*)-**71**). This piecewise procedure is also effective with primary and secondary alcohol partners in conjunction with trimethylboroxine³⁶ (**72**) or alkylboranes (**73**).

Table 2. Deoxygenative Alkylation of *gem*-Dibromoolefins^a

^aReactions performed with alcohol (1.3 equiv), *gem*-dibromoolefin (0.5 mmol, 1.0 equiv), NHC (1.4 equiv), pyridine (1.4 equiv), [Ir(ppy)₂(dtbbpy)]PF₆ (1.5 mol %), Ni(dtbbpy)Br₂ (10 mol %), quinuclidine (1.4 equiv), phthalimide (1.0 equiv), MTBE:1,2-dichlorobenzene (1:2, 0.033 M), integrated photoreactor (450 nm, m1 plate, 100% light intensity), 2 h. ^b15 mL DCB used. ^c1.75 equiv of alcohol used. ^dAssay yield (by NMR vs mesitylene as internal standard). ^e1.5 equiv of alcohol used. ^f0.45 mmol scale. ^g0.4 mmol scale. ^hTrifluorotoluene used as cosolvent with MTBE. See Supporting Information for experimental details.




Figure 2. Sequential strategies toward piecewise assembly of multisubstituted alkenes. (See the [Supporting Information](#) for experimental details.)

We next envisioned a one-flask synthesis of tetrasubstituted alkenes via sequential addition of two distinct NHC-activated alcohol partners to a reaction vessel containing the *gem*-dibromoolefin substrate. To our delight, this iterative procedure can be deployed using methyl (**74**, **75**, **76**), primary (**75**, **77**), and secondary (**74**) alcohol partners, producing tetrasubstituted alkenes in high yields with excellent cross-selectivity. Additionally, 1,4-butanediol can form a tetrasubstituted alkene in a single step (**78**), wherein the doubly NHC-activated diol can engage in coupling and subsequent 5-*exo*-trig cyclization to furnish the desired product.

Stereodefined alkenes are prevalent motifs in natural products, yet their stereoselective synthesis often relies on substrate-specific strategies tailored to the alkene target. To highlight the modularity of our approach, we envisioned its application to the synthesis of (+)-sponalisolide B, a natural product isolated from *Spongia officinalis* that exhibits quorum-sensing inhibitory activity.³⁷ Deoxygenative alkylation enables direct synthesis of (+)-sponalisolide B from simple starting materials. Moreover, by using an alternative commercially available alcohol, we synthesized a homologue (**81**) that is inaccessible via previously reported protocols (Figure 2). More broadly, this modular approach provides common alkenyl bromide intermediates en route to stereodefined alkene-containing natural products and their analogues.

A plausible mechanism for the deoxygenative bromoalkenylation of *gem*-dibromoolefins is shown in Figure 3. Following activation of the alcohol by NHC, blue light irradiation generates an excited-state iridium photocatalyst, which undergoes reductive single-electron transfer (SET) with the alcohol-NHC adduct **83** to form the reduced photocatalyst and the alcohol-NHC radical cation, **83**^{•+}. Subsequent deprotonation and rapid β -scission of **83**^{•+} generates the desired alkyl radical




Figure 3. Proposed mechanism.

(**85**[•]). Concurrently, reduced nickel(I) species **86** undergoes facile oxidative addition with the *gem*-dibromoolefin to

generate a bromovinyl nickel(III) species, **87**. Separately, the reduced iridium photocatalyst can undergo SET with a nickel(II) species (**88**) to yield nickel(I) species **89**. Comproportionation³⁵ of **89** and **87** furnishes a nickel(II) species (**90**), primed for radical capture of alkyl radical **85**[•]. Rapid reductive elimination of the resultant nickel(III) species (**91**) generates the desired deoxygenative bromoalkenylation product and reforms **86**. We propose that the high stereoselectivity observed with aldehyde-derived *gem*-dibromoolefins can be attributed to high selectivity in the oxidative addition toward the C–Br bond furthest from the alkene substituent. We further propose that phthalimide may suppress subsequent oxidative addition of the deoxygenative bromoalkenylation product to the nickel by reducing Ni(I) reactivity (see [Supporting Information](#)).

Herein we have reported the direct deoxygenative nickel metallaphotoredox-catalyzed cross-coupling of alcohols with alkenyl bromides and *gem*-dibromoolefins to generate multisubstituted alkenes and stereodefined alkenyl bromides. These reactions proceed with excellent functional group tolerance and—in the case of trisubstituted alkenes and alkenyl bromides—high stereoselectivity. This platform enables iterative, piecewise assembly of multisubstituted alkenes, providing a new pathway for the synthesis of these highly valued motifs from feedstock building blocks. We anticipate that the modularity and functional group tolerance of this platform will find broad application in the synthesis of complex alkene-containing molecules.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.6c09135>.

Additional experimental details, optimization studies, mechanistic studies, compound characterization, and spectra ([PDF](#))

■ AUTHOR INFORMATION

Corresponding Author

David W. C. MacMillan — Merck Center for Catalysis at Princeton University, Princeton, New Jersey 08544, United States; orcid.org/0000-0001-6447-0587; Email: dmacmill@princeton.edu

Authors

Eli Jones — Merck Center for Catalysis at Princeton University, Princeton, New Jersey 08544, United States

Robert T. Martin — Merck Center for Catalysis at Princeton University, Princeton, New Jersey 08544, United States; orcid.org/0000-0002-7676-2840

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacs.6c09135>

Author Contributions

[†]E.J. and R.T.M. contributed equally.

Notes

The authors declare the following competing financial interest(s): D.W.C.M. declares a competing interest with respect to the integrated photoreactor.

■ ACKNOWLEDGMENTS

Research reported in this work was supported by the National Institute of General Medical Sciences of the National Institutes of Health (R35GM134897) and the Princeton Catalysis Initiative. E.J. thanks the National Science Foundation for a graduate fellowship through the Graduate Research Fellowship Program (DGE-2444107). R.T.M. thanks the National Science Foundation for financial support through the MPS-Ascend Postdoctoral Research Fellowship (2316541). The authors thank Rebecca Lambert for assistance in preparing this manuscript and Dr. Brandon Kennedy and Lotus Separation LLC at Princeton University for help with purification. The authors further thank William Zhao and Ruizhe Chen for helpful discussions.

■ REFERENCES

- (1) Ertl, P.; Schuhmann, T. A Systematic Cheminformatics Analysis of Functional Groups Occurring in Natural Products. *J. Nat. Prod.* **2019**, *82* (5), 1258–1263.
- (2) Knochel, P.; Molander, G. A. *Additions to and Substitutions at C-C π -Bonds*; Comprehensive Organic Synthesis, Volume 4; Elsevier: Amsterdam, 2014.
- (3) Tobrman, T.; Hron, V. Trisubstituted Alkenes as Valuable Building Blocks. *Molecules* **2025**, *30* (16), 3370.
- (4) Maryanoff, B. E.; Reitz, A. B. The Wittig Olefination Reaction and Modifications Involving Phosphoryl-Stabilized Carbanions. Stereochemistry, Mechanism, and Selected Synthetic Aspects. *Chem. Rev.* **1989**, *89* (4), 863–927.
- (5) Brotons-Rufes, A.; Posada-Pérez, S.; Martínez, J. P.; Nolan, S. P.; Poater, A. Challenges in Olefin Metathesis: Past, Present and Future. *Coord. Chem. Rev.* **2025**, *542*, 216827.
- (6) Buttard, F.; Sharma, J.; Champagne, P. A. Recent Advances in the Stereoselective Synthesis of Acyclic All-Carbon Tetrasubstituted Alkenes. *Chem. Commun.* **2021**, *57* (34), 4071–4088.
- (7) Till, N. A.; Smith, R. T.; MacMillan, D. W. C. Decarboxylative Hydroalkylation of Alkynes. *J. Am. Chem. Soc.* **2018**, *140* (17), 5701–5705.
- (8) Kutateladze, D. A.; Mai, B. K.; Dong, Y.; Zhang, Y.; Liu, P.; Buchwald, S. L. Stereoselective Synthesis of Trisubstituted Alkenes via Copper Hydride-Catalyzed Alkyne Hydroalkylation. *J. Am. Chem. Soc.* **2023**, *145* (32), 17557–17563.
- (9) Ouzounthanasis, K. A.; Rizos, S. R.; Koumbis, A. E. Julia-Kocienski Olefination in the Synthesis of Trisubstituted Alkenes: Recent Progress. *Eur. J. Org. Chem.* **2023**, *26* (36), No. e202300626.
- (10) Ganiu, M. O.; Cleveland, A. H.; Paul, J. L.; Kartika, R. Triphosgene and DMAP as Mild Reagents for Chemoselective Dehydration of Tertiary Alcohols. *Org. Lett.* **2019**, *21* (14), 5611–5615.
- (11) Miyaura, N.; Ishiyama, T.; Ishikawa, M.; Suzuki, A. Palladium-Catalyzed Cross-Coupling Reactions of B-Alkyl-9-BBN or Trialkylboranes with Aryl and 1-Alkenyl Halides. *Tetrahedron Lett.* **1986**, *27* (52), 6369–6372.
- (12) Wei, L.; Popescu, M. V.; Noble, A.; Paton, R. S.; Aggarwal, V. K. Boron-Mediated Modular Assembly of Tetrasubstituted Alkenes. *Nature* **2025**, *643* (8073), 975–982.
- (13) Noble, A.; McCarver, S. J.; MacMillan, D. W. C. Merging Photoredox and Nickel Catalysis: Decarboxylative Cross-Coupling of Carboxylic Acids with Vinyl Halides. *J. Am. Chem. Soc.* **2015**, *137* (2), 624–627.
- (14) Noble, A.; MacMillan, D. W. C. Photoredox α -Vinylolation of α -Amino Acids and *N*-Aryl Amines. *J. Am. Chem. Soc.* **2014**, *136* (33), 11602–11605.
- (15) Johnson, K. A.; Biswas, S.; Weix, D. J. Cross-Electrophile Coupling of Vinyl Halides with Alkyl Halides. *Chem. - Eur. J.* **2016**, *22* (22), 7399–7402.
- (16) Pawley, S. B.; Conner, A. M.; Omer, H. M.; Watson, D. A. Development of a General Method for the Hiyama-Denmark Cross-

- Coupling of Tetrasubstituted Vinyl Silanes. *ACS Catal.* **2022**, *12* (20), 13108–13115.
- (17) Dong, Z.; MacMillan, D. W. C. Metallaphotoredox-Enabled Deoxygenative Arylation of Alcohols. *Nature* **2021**, *598* (7881), 451–456.
- (18) Gould, C. A.; Pace, A. L.; MacMillan, D. W. C. Rapid and Modular Access to Quaternary Carbons from Tertiary Alcohols via Bimolecular Homolytic Substitution. *J. Am. Chem. Soc.* **2023**, *145* (30), 16330–16336.
- (19) Lin, E.; Wang, J. Z.; Mao, E.; Tsang, S.; Carsch, K. M.; Prieto Kullmer, C. N.; McNamee, R. E.; Long, J. R.; Le, C. “Chip”; MacMillan, D. W. C. Aryl Acid-Alcohol Cross-Coupling: C(sp³)-C(sp²) Bond Formation from Nontraditional Precursors. *J. Am. Chem. Soc.* **2025**, *147* (18), 14905–14914.
- (20) Lyon, W. L.; MacMillan, D. W. C. Expedient Access to Underexplored Chemical Space: Deoxygenative C(sp³)-C(sp³) Cross-Coupling. *J. Am. Chem. Soc.* **2023**, *145* (14), 7736–7742.
- (21) Großkopf, J.; Gopatta, C.; Martin, R. T.; Haseloer, A.; MacMillan, D. W. C. Generalizing Arene C-H Alkylations by Radical-Radical Cross-Coupling. *Nature* **2025**, *641* (8061), 112–121.
- (22) Gould, C. A.; Pace, A. L.; MacMillan, D. W. C. Rapid and Modular Access to Quaternary Carbons from Tertiary Alcohols via Bimolecular Homolytic Substitution. *J. Am. Chem. Soc.* **2023**, *145* (30), 16330–16336.
- (23) Ertl, P.; Schuhmann, T. A Systematic Cheminformatics Analysis of Functional Groups Occurring in Natural Products. *J. Nat. Prod.* **2019**, *82* (5), 1258–1263.
- (24) Spaggiari, A.; Vaccari, D.; Davoli, P.; Torre, G.; Prati, F. A Mild Synthesis of Vinyl Halides and *gem*-Dihalides Using Triphenyl Phosphite-Halogen-Based Reagents. *J. Org. Chem.* **2007**, *72* (6), 2216–2219.
- (25) Uehling, M. R.; Rucker, R. P.; Lalic, G. Catalytic Anti-Markovnikov Hydrobromination of Alkynes. *J. Am. Chem. Soc.* **2014**, *136* (24), 8799–8803.
- (26) Imazaki, Y.; Shirakawa, E.; Ueno, R.; Hayashi, T. Ruthenium-Catalyzed Transformation of Aryl and Alkenyl Triflates to Halides. *J. Am. Chem. Soc.* **2012**, *134* (36), 14760–14763.
- (27) Das, J. P.; Roy, S. Catalytic Hunsdiecker Reaction of α,β -Unsaturated Carboxylic Acids: How Efficient Is the Catalyst? *J. Org. Chem.* **2002**, *67* (22), 7861–7864.
- (28) Desai, N. B.; McKelvie, N.; Ramirez, F. A New Synthesis of 1,1-Dibromoolefins *via* Phosphine-Dibromomethylenes. the Reaction of Triphenylphosphine with Carbon Tetrabromide. *J. Am. Chem. Soc.* **1962**, *84* (9), 1745–1747.
- (29) Konowalchuk, D. J.; Hall, D. G. Solvent-Controlled Monoselective Suzuki-Miyaura Cross-Coupling of Symmetrical 1,1-Dichloroalkenes: Scope and Origin of Selectivity. *ACS Catal.* **2026**, *16* (3), 2698–2710.
- (30) Novoa, A.; Pellegrini-Moise, N.; Bourg, S.; Thoret, S.; Dubois, J.; Aubert, G.; Cresteil, T.; Chapleur, Y. Design, Synthesis and Antiproliferative Activities of Biaryllolefins Based on Polyhydroxylated and Carbohydrate Scaffolds. *Eur. J. Med. Chem.* **2011**, *46* (9), 3570–3580.
- (31) Shen, W. A Modified Suzuki Reaction of 1,1-Dibromo-1-Alkenes. *Synlett* **2000**, *2000* (5), 0737–0739.
- (32) Bauer, A.; Miller, M. W.; Vice, S. F.; McCombie, S. W. Suzuki Arylation of 1,1-Dibromo-1-Alkenes: Synthesis of Tetra-Substituted Alkenes. *Synlett* **2001**, *2001* (02), 0254–0256.
- (33) Renzi, P.; Lanfranco, A.; Pazderová, L.; Rusconi, M.; Scarfiello, J.; Deagostino, A. Synthetic Methodologies to Access Skipped Dienes: A Focus on the Catalytic Systems. *Org. Biomol. Chem.* **2025**, *23* (25), 6052–6099.
- (34) Prieto Kullmer, C. N.; Kautzky, J. A.; Krska, S. W.; Nowak, T.; Dreher, S. D.; MacMillan, D. W. C. Accelerating reaction generality and mechanistic insight through additive mapping. *Science* **2022**, *376* (6592), 532–539.
- (35) Raab, T. J.; Doyle, A. G. Reactivity Studies of Bipyridine-Ligated Nickel(I) and Nickel(0) Complexes Inform the Mechanism

in Modern Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2025**, *147* (37), 33991–34000.

(36) Gray, M.; Andrews, I. P.; Hook, D. F.; Kitteringham, J.; Voyle, M. Practical Methylation of Aryl Halides by Suzuki-Miyaura Coupling. *Tetrahedron Lett.* **2000**, *41* (32), 6237–6240.

(37) Sun, D.-Y.; Han, G.-Y.; Yang, N.-N.; Lan, L.-F.; Li, X.-W.; Guo, Y.-W. Racemic Trinorsesquiterpenoids from the Beihai Sponge *Spongia officinalis*: Structure and Biomimetic Total Synthesis. *Org. Chem. Front.* **2018**, *5* (6), 1022–1027.