## **Accelerated Article Preview**

# Alkene dialkylation by triple radical sorting

Received: 20 November 2023

Accepted: 5 February 2024

Accelerated Article Preview

Cite this article as: Wang, J. Z. et al. Alkene dialkylation by triple radical sorting. *Nature* https://doi.org/10.1038/s41586-024-07165-x (2024) Johnny Z. Wang, William L. Lyon & David W. C. MacMillan

This is a PDF file of a peer-reviewed paper that has been accepted for publication. Although unedited, the content has been subjected to preliminary formatting. Nature is providing this early version of the typeset paper as a service to our authors and readers. The text and figures will undergo copyediting and a proof review before the paper is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers apply.

### Alkene dialkylation by triple radical sorting

Johnny Z. Wang,<sup>1,2</sup> William L. Lyon,<sup>1,2</sup> David W. C. MacMillan<sup>1\*</sup>

The development of bimolecular homolytic substitution ( $S_{H2}$ ) catalysis has expanded cross-coupling chemistries by enabling the selective combination of any primary radical with any secondary or tertiary radical via a radical sorting mechanism<sup>1-8</sup>. Biomimetic<sup>9,10</sup> S<sub>H2</sub> catalysis can be used to merge common feedstock chemicals—such as alcohols, acids, and halides—in various permutations for the construction of a single  $C(sp^3)-C(sp^3)$  bond. The ability to sort these two distinct radicals across commercially available alkenes in a three-component manner would enable the simultaneous construction of two  $C(sp^3)-C(sp^3)$ bonds, greatly accelerating access to complex molecules and drug-like chemical space<sup>11</sup>. However, the simultaneous *in situ* formation of electrophilic and primary nucleophilic radicals in the presence of unactivated alkenes is problematic, typically leading to statistical radical recombination, hydrogen atom transfer, disproportionation, and other deleterious pathways<sup>12,13</sup>. Herein, we report the use of bimolecular homolytic substitution catalysis to sort an electrophilic radical and a nucleophilic radical across an unactivated alkene. This reaction involves the *in situ* formation of three distinct radical species, which are then differentiated by size and electronics, allowing for regioselective formation of the desired dialkylated products. This work accelerates access to pharmaceutically relevant C(sp<sup>3</sup>)-rich molecules and defines a distinct mechanistic approach for alkene dialkylation.

A key goal of organic chemistry is the development of new methods for the rapid synthesis of  $C(sp^3)-C(sp^3)$  bonds en route to threedimensional drug-like molecules.<sup>14-16</sup>. Traditional cross-coupling paradigms rely on oxidative addition, transmetallation, and reductive elimination mechanistic steps that limit the pool of potential coupling partners. By contrast, bimolecular homolytic substitution (S<sub>H</sub>2) catalysis couples primary radicals with secondary or tertiary radicals through a radical sorting mechanism based on carbon–metal bond strength<sup>1-8,17</sup>. As this unique radical sorting mechanism is functional group-agnostic, abundant radical precursors, such as alcohols, acids, and halides, can be coupled in any desired combination to generate complex products from simple feedstock chemicals<sup>1-8</sup> (Fig 1a). This novel approach greatly expands access to  $C(sp^3)$ -rich chemical space and enables formation of otherwise elusive all- $C(sp^3)$  quaternary centers<sup>18</sup>. Although radical-sorting S<sub>H</sub>2 catalysis has been shown to construct a single  $C(sp^3)$  bond from a variety of radical precursors, the more challenging three-component radical sorting mechanism has yet to be demonstrated.

A one-step protocol for the regioselective dialkylation of unactivated alkenes is highly desirable<sup>11</sup>. Alkenes are widely available, and the simultaneous construction of two  $C(sp^3)$ – $C(sp^3)$  bonds across an alkene would greatly accelerate access to therapeutically advantageous  $C(sp^3)$ -rich small molecules<sup>14</sup>. Due to the propensity of alkyl-metal complexes to undergo  $\beta$ –H elimination<sup>19-21</sup>, existing methods for alkene dialkylation remain greatly limited, relying either on auxiliary functional groups to direct dialkylation<sup>22-24</sup> or the presence of specific ground-state radical traps<sup>25</sup>. A general method in which two distinct radicals are formed and regioselectively added across any unactivated alkene represents an ideal approach.

We envisioned a catalytic alkene dialkylation platform commencing with addition of an electrophilic alkyl radical—such as trifluoromethyl or difluoroacyl radical—into an alkene. Subsequent radical-radical recombination of the resultant radical species with a nucleophilic alkyl radical would yield the dialkyl adduct in a single transformation (Fig. 1b). The proposed dialkylation would be expected to proceed with good regioselectivity and enable installation of electron-poor alkyl groups that are typically not compatible with nickel catalysis<sup>26</sup>. Through the course of the reaction, three distinct radicals would be formed and must be efficiently sorted. Unsurprisingly, the envisioned transformation does not proceed in the absence of any sorting catalysts; instead, deleterious pathways, including disproportionation, alkyl-alkyl dimerization, and hydrogen atom transfer (HAT) predominate <sup>12,13</sup>. We postulated that an appropriate S<sub>H</sub>2 catalyst, which has been shown to facilitate outer-sphere C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond formation, might be used to sort these three simultaneously generated radicals toward productive alkene dialkylation. For the hypothesized radical sorting to be operative, the catalyst used must be capable of preferentially binding primary alkyl radicals over high-energy electrophilic alkyl radicals and secondary or tertiary radicals, while still being capable of performing outer-sphere bimolecular homolytic substitution for C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond formation. Herein, we disclose a general strategy for the dialkylation of alkenes through the simultaneous generation of electrophilic alkyl radicals and primary nucleophilic radicals in the presence of unactivated alkenes (Fig. 1c). The functional group-agnostic nature of S<sub>H</sub>2 catalysis permits the use of commercially available primary alcohols and electron-poor alkyl chlorides as radical precursors, giving potential access to  $2 \times 10^{15}$  C(sp<sup>3</sup>)-rich dialkylated products<sup>27</sup>.

#### Mechanism

We envisioned that the alkene dialkylation would proceed via the mechanism outlined in Figure 2. Condensation of primary alcohol 1 onto a benzoxazolium salt (NHC) forms adduct 2 in situ<sup>28</sup>. Meanwhile, blue light excitation of photocatalyst 3 accesses a long-lived, triplet excited state 4 ( $E_{1/2}^{red}$  [\*Ir<sup>III</sup>/Ir<sup>II</sup>]=+1.21 vs. saturated calomel electrode (SCE) in MeCN)<sup>29</sup>. Stern-Volmer analysis (See SI pages S32–S33) suggests that 4 undergoes reductive quenching with 2. Subsequent deprotonation and facile  $\beta$ -scission, provides the desired primary alkyl radical (6) and a benign aromatized byproduct. The primary alkyl radical can then be captured by high valent nickel S<sub>H</sub>2 catalyst 7, producing nickel–alkyl complex 8<sup>5</sup>. To close the photocatalytic cycle, reduced-state Ir<sup>II</sup> (9) is capable of

51

1

<sup>52</sup> 53 54

<sup>&</sup>lt;sup>1</sup>Merck Center for Catalysis at Princeton University, Princeton, NJ 08544, USA. <sup>2</sup>These authors contributed equally: Johnny Z. Wang, William L. Lyon. \*email: dmacmill@princeton.edu

reducing an α-acyl alkyl chloride **10** (or dMesSCF<sub>3</sub>(OTf)) to produce an electrophilic carbon-centered radical (**11**) that can add into an unactivated alkene (**12**). Radical probes (See SI pages **S28–S31**) support the formation of tertiary radical (**13**) which is capable of being further functionalized<sup>30-32</sup>. This nucleophilic tertiary radical can undergo a S<sub>H</sub>2 reaction with **8**, thereby regenerating **7** and forming the desired dialkylated product (**14**) (see SI for select mechanistic experiments). Key to the success of this reaction is the radical sorting of the many transient radicals, both electronically, through addition to the alkene, and sterically, through binding to a high valent nickel complex. We envisioned that this novel mechanistic paradigm might provide a general, modular strategy for the dialkylation of unactivated olefins.

#### Alkene scope

We first sought to interrogate the scope of the alkene coupling partner (Fig 3). We selected trifluoromethyl and difluoroacetamide radicals as the electrophilic alkyl radical partners, as both groups are important in drug discovery<sup>33, 34</sup>. For the primary radical partner, we opted to use the methyl radical, both for its ability to favorably influence the properties of drugs (termed the magic methyl effect) and due to the challenge often posed by its incorporation into complex molecules<sup>35</sup>. Gratifyingly, the coupling reaction proceeded efficiently across a wide range of alkenes. Unactivated terminal alkenes with relatively low  $\pi$ -nucleophilicity<sup>36</sup> were dialkylated in good yields, tolerating protic functionality (**15**, **16**) as well as homolytically labile allyl–benzylic C–H bonds (**17**, **18**). The coordinatively saturated S<sub>H</sub>2 catalyst is incapable of oxidative addition; as such, aryl and alkyl halide-containing vinyl ethers and enamides were dialkylated in good yield (**19–21**). The generation of quaternary centers, a longstanding challenge in organic synthesis, can be achieved from 4, 5, 6, and 7-member rings as well as from acyclic 1,1-disubstituted alkenes (**22–27**). Notably, tertiary boronic esters (**28**), ethers (**29**), and ureas (**30**) were effectively formed under our reaction conditions. Moreover, a range of 1,2-disubstituted and trisubstituted alkenes were competent substrates (**31–34**). As a demonstration of the mild and robust nature of the reaction conditions, we successfully dialkylated several complex bioactive molecules. Specifically, efinconazole (**35**), paroxetine derivative (**36**), vinclozolin (**37**), retapamulin (**38**), quinine (**39**), and ataluren derivative (**40**) were dialkylated in good yields, showcasing the ability of the reaction to tolerate tertiary amines, alcohols, sulfides, quinuclidine, and oxidative addition-prone oxadiazole functionality. These results suggest that the protocol should be applicable to the late-stage dialkylation of alkenes.

#### Alkyl chloride scope

We next turned our attention to exploring the scope of the electrophilic radical partner (Fig 4). Alkyl chlorides were selected as radical precursors for their ease of synthesis, commercial availability, and enhanced stability over their bromide counterparts. We found that  $\alpha$ -acyl radicals spanning a range of electrophilicity profiles, from  $\alpha$ -ester to  $\alpha$ -difluoroester radicals (41–43), reacted in good yields. Both acetamide (44), a ubiquitous moiety in drugs, and substituted difluoroacetamides (45), which are particularly important in fragment-based drug discovery, could be incorporated through the dialkylation protocol. Synthetically useful Horner-Wadsworth-Emmons reagents (46) and  $\alpha$ -acyl chlorides (47) were prepared in good yields and offer the potential for further elaboration. In addition to  $\alpha$ -acyl radicals, we found numerous other electrophilic carbon-centered radicals to be viable electrophilic radical partners, including  $\alpha$ -difluorosulfonyl (48), difluorobenzyl (49), and  $\alpha$ -nitrile (50) radicals.

#### Primary alcohol scope

Finally, we explored the scope of the primary alcohol coupling partner. As shown in Figure 4, a broad scope of pharmaceutically relevant alkyl fragments were incorporated through our protocol. Coupling with commercially available <sup>13</sup>C methanol served to install an isotopically labelled methyl group at the quaternary center (**51**). Moreover, a range of alcohols were found to be effective alkyl coupling partners, including ether alcohol (**52**), threonine (**53**), serine (**54**), guaifenesin (**55**), and other diols (**56**, **57**). Reaction of diol substrates proceeds with full regioselectivity for the primary alcohol. These complex, coupled products (**53-57**) bear a free hydroxyl group that can be subjected to further elaboration via NHC activation. Moreover, 1,1-disubstituted alkenes in 4-membered rings (**58**) or acyclic substrates (**59**) were observed to undergo efficient alkylation with Cbz-glycinol. Notably, 1,2-disubstituted alkenes (**60**) were also readily dialkylated, providing an orthogonally protected morpholine scaffold. The complexity-building potential of this protocol was demonstrated through elaboration of dialkylated products into complex C(sp<sup>3</sup>) rich frameworks (**61**, **63**). The tertiary alcohol of **61** was activated by NHC, and subsequent benzylation<sup>8</sup> proceeded efficiently to generate a second quaternary center (**62**). In the case of **63**, the tertiary alcohol served as a radical precursor en route to alkylation with dehydroalanine<sup>37</sup> to yield complex scaffold, **64**.

#### Conclusion

Key to the strategy described herein is an outer-sphere  $C(sp^3)$ – $C(sp^3)$  bond formation capable of forming quaternary centers. A wide range of unactivated alkenes were dialkylated, including tertiary amines, alcohols, aryl halides, and other reactive functionalities. Several examples of both the electrophilic radical and primary radical partners containing sites for further elaboration were demonstrated in good yield. Modulation of all three reaction components should allow for the rapid synthesis of  $C(sp^3)$ -rich small molecule libraries. Furthermore, the described approach provides a framework for future developments in  $C(sp^3)$ – $C(sp^3)$  bond-forming alkene difunctionalization.

#### References

117

198

- Liu, W., Lavagnino, M. N., Gould, C. A., Alcazar, J. & MacMillan, D. W. C. A biomimetic SH2 cross-coupling mechanism for quaternary sp<sup>3</sup>-carbon formation. Science. 1. 6572. 1258-1263 (2021)
- Tsymbal, A. V., Bizzini, L. D. & MacMillan, D. W. C. Nickel catalysis vis SH2 homolytic substitution: the double decarboxylative cross-coupling of aliphatic acids. J. Am. 2. Chem. Soc. 144, 21278-21286 (2022)
- 3. Mao, E. & MacMillan D. W. C. Late-stage C(sp3)-H methylation of drug molecules. J. Am. Chem. Soc. 145, 2787-2793 (2023)

Sakai, H. A. & MacMillan, D. W. C., Nontraditional fragment coupling of alcohol and carboxylic acids: C(sp3)-C(sp3) cross-coupling via radical sorting. J. Am. Chem. 4. Soc. 144, 6185-6192, (2022)

5 Gould, C. A., Pace, A. L. & MacMillan, D. W. C., Rapid and modular access to quaternary carbons from tertiary alcohols via bimolecular homolytic substitution. J. Am. Chem. Soc. 145, 16330-16336 (2023)

Gan, X., Kotesova, S., Castanedo, A., Green, S. A., Moller, S. L. B. & Shenvi, R. A., Iron-catalyzed hydrobenzylation: Stereoselective synthesis of (-)-Eugenial C. J. Am. 6. Chem. Soc. 145, 15714-15720 (2023)

- 7. Ga., X., et. al. Carbon guaternization of redox active esters and olefins via decarboxylative coupling. ChemRxiv. November 17, 2023. DOI: 10.26434/chemrxiv-2023-7vb8x (Accessed 2023-11-19)
- 8. Lovett, V. P., Kong, L., Gan, X. & Shenvi, R. A. Single catalyst double outer-sphere alkene cross-coupling. ChemRxiv. November 17, 2023. DOI: 10.26434/chemrxiv-9c2 (Accessed 2023-11-19)

Zhang, Q., van der Donk, W. A. & Liu, W. Radical-mediated enzymatic methylation: A tale of two SAMS. Acc. Chem. Res. 45, 555-564 (2012)

- Bauerle, M. R., Schwalm, E. L. & Booker, S. J. Mechanistic diversity of radical S-adenomethionine (SAM)-dependent methylation. J. Biol. Chem. 290, 3995–4002 (2015) 10 Qi, X. & Diao, T. Nickel-catalyzed dicarbofunctionalization of alkenes. ACS Catal. 10, 8542-8556 (2020) 11
- 12
- Gibian, M. J. & Corley, R. C. Organic radical-radical reactions. disproportionations vs. combination. *Chem. Rev.* 73, 441–465 (1973) Leifert, D. & Studer, A. The persistent radical effect in organic synthesis. *Angew. Chem. Int. Ed.* 59, 74–108 (2020) 13
- 14.
- Lovering F. Escape from fatland 2: complexity and promiscuity. *Med. Chem. Commun.* **4**, 515–519 (2013) Choi, J. & Fu, G. C., Transition metal-catalyzed alkyl-alkyl bond formation: Another dimension in cross-coupling chemistry. *Science.* **356**, eaaf7230 (2017) 15.
- Dombrowski, A. W., et. al. Expanding the medicinal chemist toolbox: Comparing seven C(sp<sup>2</sup>)-C(sp<sup>3</sup>) cross-coupling methods by library synthesis. ACS Med. Chem. 16. Lett. 11, 597-604 (2020)
- 17 Simoes, J. A. M. & Beauchamp, J. L. Transition metal-hydrogen and metal-carbon bond strengths: the keys to catalysis. Chem. Rev. 90, 629–688 (1990)
- Quasdorf, K. W. & Overman, L. E. Catalytic enantioselective synthesis of quaternary carbon stereocenters. Nature, 516, 181–191 (2014) 18
- Dhungana R. K., Sapkota, R. R., Wickham, L. M., Niroula, D. & Giri, R. Ni-Catalyzed regioselective 1,2-dialkylation of alkenes enabled by the formation of two C(sp<sup>3</sup>)-19 C(sp<sup>3</sup>) bonds. J. Am. Chem. Soc. 142, 20930-20936 (2020)
- Jana, R., Pathak, T. P. & Sigman, M. S. Advances in transition metal (Pd, Ni, Fe)-catalyzed cross-coupling reactions using alkyl-organometallics as reaction partners. 20 Chem. Rev. 111, 1417–1492 (2011)
- Frisch, A. C. & Beller, M. Catalysts for cross-coupling reactions with non-activated alkyl halides. Angew. Chem. Int. Ed. 44, 674-688 (2005) 21
- Zhang, J. X. & Shu, W. Ni-Catalyzed reductive 1,2-cross-dialkylation of unactivated alkenes with two alkyl bromides. Org. Lett. 24, 3844–3849 (2022)
- 23. Dorsea, J., Puyl, V. A., Tran, V. A., Liu, M. & Engle, K. M., Directed nickel-catalyzed 1,2-dialkylation of alkenyl carbonyl compounds. Chem. Sci. 9, 5278–5283 (2018). Yang, T., Jiang, Y., Luo, Y., Lim, J. H., Lan, Y. & Koh, M. J. Chemoselective union of olefins, organohalides, and redox-active esters enables regioselective alkene 24
- dialkylation. J. Am. Chem. Soc. 142, 21410-21419 (2020) Rao C. Q., Zhang, T. Z., Liu, H. C. & Huan, H. M. Double alkyl-alkyl bond construction across alkenes enabled by nickel electron-shuttle catalysis, Nat. Catal. (2023) 25
- Le, C., Chen, T. Q., Liang, Y., Zhang, P. MacMillan D. W. C. A radical approach to the copper oxidative addition problem: trifluoromethylation of bromoarenes. Science, 26 360. 1010-1014 (2018)
- Reaxys search from October 2023 of commercially abailable fragments: 1º alcohols (261,696), α-acyl chlorides (29,000), alkenes (268553). 27
- Dong, Z. & MacMillan, D. W. C., Metallaphotoredox-enabled deoxygenative arylation of alcohols. Nature. 598, 451–456 (2021). 28
- Lowry, M. S., et. al. Single-layer electroluminescent devices and phtoinduced hydrogen production form an ionic iridium(III) complex. Chem. Mater. 17, 5712–5719, 29 (2005)
- Kharasch, M. S., Skell, P. S. & Fisher, P., Reactions of atoms and free radicals in solution. XII. The addition of bromo esters to olefins. J. Am. Chem. Soc. 70, 1055-30 1059 (1948)
- Bian, K. J., Nemoto Jr, D., Kao, S. C., He, Y., Li, Y., Wang, X. S. & West, J. G. Modular difunctionalization of unactivated alkenes through bio-inspired radical ligand 31 transfer catalysis. J. Am. Chem. Soc. 144, 11810-11821 (2022)
- 32 Patra, S., Giri, R. & Katayev, D. Nitrative difunctionalization of alkenes via cobalt-mediated radical ligand transfer and radical-polar crossover photoredox catalysis. ACS Catal. 13, 16136-16147 (2023)
- Nair, A. S., et. al. FDA-approved trifluoromethyl group-containing drugs: A review of 20 years. Processes. 10, 2054–2078 (2022)
- Vaas, S., et. al. Principles and applications of CF<sub>2</sub>X moieties as unconventional halogen bond donors in medicinal chemistry, chemical biology, and drug discovery. J. 34 Med. Chem. 66, 10202-10225 (2023)
- Schönherr, H. & Cernak, T. Profound methyl effects in drug discovery and a call for new C-H methylation reactions. Angew. Chem. Int. Ed. 52, 12256–12267 (2013)
- Mayr, H., Kempf, B. & Ofial, A. R., π-Nucleophilicity in carbon-carbon bond-forming reactions. Acc. Chem. Res. 36, 66-77 (2003) 36
- 37 Wang, J. Z., Sakai, H. A. & MacMillan D. W. C., Alcohols as alkylating agents: Photoredox-catalyzed conjugate alkylation via in situ deoxygenation. Angew. Chem. Int. Ed. 61, e202207150 (2022)

Data Availability: The data supporting the findings of this study are available within the paper and its Supplementary Information.

Supplementary Information is linked to the online version of the paper at www.nature.com/nature

Acknowledgements We thank Nathan W. Dow and Colin A. Gould for helpful scientific discussion. Research reported in this work was supported by the NIH National Institute of General Medical Sciences (R35 GM134897-04) and kind gifts from Merck, Pfizer, Janssen, Bristol-Myers Squibb, Genentech, and Genmab. J.Z.W. and W.L.L. acknowledge Princeton University, E. Taylor, and the Taylor family for an Edward C. Taylor Fellowship. W.L.L. acknowledges the National Science Foundation for a predoctoral fellowship (Award DGE-2039656). The authors thank Rebecca Lambert for assistance in preparing this manuscript.

Author Contributions J.Z.W., W.L.L., and D.W.C.M. designed the experiments. J.Z.W., and W.L.L. performed and analyzed the experiments. J.Z.W., and D.W.C.M. prepared this manuscript.

Competing Interests D.W.C.M. declares a competing financial interest with respect to the Penn PhD Integrated Photoreactor, which is used to irradiate reactions in this work. The remaining authors declare no competing interests.

Author Information Reprints and permissions information is available at www.nature.com/reprints. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to D.W.C.M. (dmacmill@princeton.edu).

Figure 1 Radical sorting-enabled alkene dialkylation. a, Bimolecular homolytic substitution (S<sub>H</sub>2) radical sorting enables the use of any functional group in any combination for C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond formation. **b**, Three-component radical sorting enables alkene dialkylation. **c**, This work: alkene dialkylation of unactivated alkenes using primary alcohols and α-acyl chlorides as radical precursors. Boc, tert-butylcarbonyl; Me, methyl; Cbz, carbobenzyloxy; Ph, phenyl; Bn, Benzyl; NHC, Nheterocyclic carbene; Ni, nickel; PC, photocatalyst; Ar, aryl.

**Figure 2 Proposed mechanism of alkene dialkylation** Alk, Alkyl; *t*-Bu, *tert*-butyl; EWG, electron withdrawing group; dMesCF<sub>3</sub>(OTf), Dimesityl(trifluoromethyl)sulfonium trifluoromethanesulfonate; L, ligand.

**Figure 3 Alkene scope** <sup>a</sup>MeOH (2 equiv.), NHC-1 (2.1 equiv.), Pyridine (2.1 equiv.), Ni(acac)<sub>2</sub> (15 mol%), KTp\* (15 mol%), (Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy))PF<sub>6</sub> (1 mol%), dMesSCF<sub>3</sub>(OTf) (2 equiv.), alkene (0.50 mmol), CsOAc (2.5 equiv.), TBME/tAmOH (1:1, 0.05 M), IPR(2% light intensity, 12 h). <sup>b</sup>MeOH (2 equiv.), NHC-1 (2.1 equiv.), Pyridine (2.1 equiv.), Ni(acac)<sub>2</sub> (25 mol%), 4CzPN (1 mol%), difluorochloroacetamide (2 equiv.), alkene (0.50 mmol), CsOAc (2.5 equiv.), TBME/MeCN (1:1, 0.05 M), IPR (50% light intensity, 12 h). Isolated yields. Ir, iridium; Ni(acac)(Tp\*), Nickel(acetylacetone)tris(3,5-dimetyl-1-pyrazolyl)borate; Ir-1, Ir(dFCF<sub>3</sub>ppy)<sub>2</sub>(dtbbpy)PF<sub>6</sub>; TBS, *tert*-butyldimethylsilyl; Ln, ligand; Ni(acac)<sub>2</sub>, nickel(II) bis(acetylacetone); KTp\*, potassium tris(3,5-dimetyl-1-pyrazolyl)borate; 4CzPN, 3,4,5,6-tetra(9H-carbazol-9-yl)phthalonitrile; IPR, integrated photoreactor.

**Figure 4 Scope of electrophilic and nucleophilic radicals** <sup>a</sup>MeOH (2.0 equiv.), NHC-1 (2.1 equiv.), Pyridine (2.1 equiv.), Ni(acac)<sub>2</sub> (25 mol%), 4CzPN (1 mol%), alkyl chloride (2 equiv.), alkene (0.50 mmol), CsOAc (2.5 equiv.), TBME/MeCN (1:1, 0.05 M). <sup>b</sup>primary alcohol (2.5 equiv.), NHC-*p*CF<sub>3</sub> (2.6 equiv.), Pyridine (2.6 equiv.), Ni(TMHD)<sub>2</sub> (25 mol%), 4CzPN (1 mol%), alkyl chloride (2.5 equiv.), alkene (0.50 mmol), CsOAc (3.0 equiv.), TBACI (0.6 equiv.) TBME/tAmOH (1:1, 0.05 M), IPR (5% intensity, 16 h). <sup>c</sup>See supplementary information for experimental details. <sup>d</sup>Yield determined by <sup>19</sup>F NMR analysis with 1,4-difluorobenzene as an internal standard. <sup>e</sup>Utilizing 3-(trifluoromethyl)benzyl bromide as a cross-coupling partner; see supplementary information for experimental details. All yields are isolated unless otherwise noted. Et, ethyl, Ni(TMHD)<sub>2</sub>, nickel(II) bis(2,2,6,6-tetramethyl-3,5-heptanedionate).







