

Bioinspired Supercharging of Photoredox Catalysis for Applications in Energy and Chemical Manufacturing

Agustin Millet, Paul T. Cesana, Kassandra Sedillo, Matthew J. Bird, Gabriela S. Schlau-Cohen, Abigail G. Doyle, David W. C. MacMillan,* and Gregory D. Scholes*



CONSPECTUS: For more than a decade, photoredox catalysis has been demonstrating that when photoactive catalysts are irradiated with visible light, reactions occur under milder, cheaper, and environmentally friendlier conditions. Furthermore, this methodology allows for the activation of abundant chemicals into valuable products through novel mechanisms that are otherwise inaccessible. The photoredox approach, however, has been primarily used for pharmaceutical applications, where its implementation has been highly effective, but typically with a more rudimentary understanding of the mechanisms involved in these transformations. From a global perspective, the manufacture of everyday chemicals by the chemical industry as a whole currently accounts for 10% of total global energy consumption and generates 7% of the world's greenhouse gases annually. In this context, the Bio-Inspired Light-Escalated Chemistry (BioLEC) Energy Frontier Research Center (EFRC) was founded to supercharge the photoredox approach for applications in chemical manufacturing aimed at reducing its energy consumption and emissions burden, by using bioinspired schemes to harvest multiple electrons to drive endothermically uphill chemical reactions. The Center comprises a diverse group of researchers with expertise that includes synthetic chemistry, biophysics, physical chemistry, and engineering. The team works together to gain a deeper understanding of the mechanistic details of photoredox reactions while amplifying the applications of these light-driven methodologies.

In this Account, we review some of the major advances in understanding, approach, and applicability made possible by this collaborative Center. Combining sophisticated spectroscopic tools and photophysics tactics with enhanced photoredox reactions has led to the development of novel techniques and reactivities that greatly expand the field and its capabilities. The Account is intended to highlight how the interplay between disciplines can have a major impact and facilitate the advance of the field. For example, techniques such as time-resolved dielectric loss (TRDL) and pulse radiolysis are providing mechanistic insights not previously available. Hypothesis-driven photocatalyst design thus led to broadening of the scope of several existing transformations. Moreover, bioconjugation approaches and the implementation of triplet—triplet annihilation mechanisms created new avenues for the exploration of reactivities. Lastly, our multidisciplinary approach to tackling real-world problems has inspired the development of efficient methods for the depolymerization of lignin and artificial polymers.

KEY REFERENCES

 Earley, J. D.; Zieleniewska, A.; Ripberger, H. H.; Shin, N. Y.; Lazorski, M. S.; Mast, Z. J.; Sayre, H.; Knowles, R. R.; McCusker, J. K.; Scholes, G. D.; Reid, O. G.; Rumbles, G. Ion Pair Reorganization Regulates Reactivity in PhotoReceived: February 8, 2022





Figure 1. Examples of BioLEC's contributions to the advancement of photoredox catalysis. From left to right: use of spectroscopic techniques as tools to increase fundamental understandings, discovery of new reactivities guided by mechanistic insights, strategies to circumvent nonspecific absorption issues, and depolymerization of native lignin and hydroxylated polymers via photoredox catalysis.

redox Catalysts.. Nat. Chem. **2022**, 10.1038/s41557-022-00911-6.¹ This paper describes the application of timeresolved dielectric loss spectroscopy to probe photoinduced dipole changes in an Ir photocatalyst to characterize the role of its counterion on excited state reactivity.

- Sayre, H.; Ripberger, H. H.; Odella, E.; Zieleniewska, A.; Heredia, D. A.; Rumbles, G.; Scholes, G. D.; Moore, T. A.; Moore, A. L.; Knowles, R. R. PCET-Based Ligand Limits Charge Recombination with an Ir(III) Photoredox Catalyst. J. Am. Chem. Soc. 2021, 143, 13034–13043.² This paper illustrates the design, synthesis, and characterization of a PCET-ligand that slows intramolecular charge recombination limitations when installed in an Ir photocatalyst for the reductive elimination of redox-active esters.
- Cesana, P. T.; Li, B. X.; Shepard, S. G.; Ting, S. I.; Hart, S. M.; Olson, C. M.; Martinez Alvarado, J. I.; Son, M.; Steiman, T. J.; Castellano, F. N.; Doyle, A. G.; MacMillan, D. W. C.; Schlau-Cohen, G. S. A Biohybrid Strategy for Enabling Photoredox Catalysis with Low-Energy Light. Chem2022, 8, 174–185..³ This paper describes the construction, characterization, and synthetic demonstration of a photosynthetic biohybrid consisting of a light-harvesting protein conjugated to a transition-metal photocatalyst to broaden absorption and enable low-energy excitation.
- Nguyen, S. T.; McLoughlin, E. A.; Cox, J. H.; Fors, B. P.; Knowles, R. R. Depolymerization of Hydroxylated Polymers via Light-Driven C-C Bond Cleavage. J. Am. Chem. Soc. 2021, 143, 12268–12277.⁴ This paper describes a photocatalytic approach for the depolymerization of hydroxylated polymers via a PCET step that leads to C-C bond cleavage and generation of well-defined products that can be functionalized and repolymerized to enter a closed recycling loop.

1. INTRODUCTION

The search for more efficient, cost-effective, and green manufacturing technologies is of paramount importance as global energy demands continue to increase. Chemical manufacturing alone accounts for 10% of total global energy consumption and is responsible for 7% of the world's greenhouse gas emissions.⁵ Fossil fuels remain the most widely employed source of energy to generate the high temperatures and pressures required to manufacture desired products. Therefore, the use of carbon-neutral sources of energy could have a tremendous impact at the global scale in energy-saving strategies and in reducing the chemical manufacturing industry's carbon footprint.⁶ Fossil fuels are often also the precursors to starting materials for the production of everyday chemicals. Transformation of abundant biomass to feedstock chemicals could alleviate the need for nonrenewable sources.

Solar energy is an attractive alternative to fossil fuels, as it is carbon-neutral and enough solar energy to power humanity for an entire year reaches the earth every hour.⁷ The key setback is not the lack of energy but the means to utilize and convert this power toward useful pathways. Over the last few decades, scientists have been improving the viability and energy conversion efficiency (averaging around 30%)⁸ of photovoltaic cells, which can provide electrical power to homes and businesses. Solar energy can similarly be converted to chemical energy when used to excite light-absorbing catalysts and take the place of thermal energy in chemical syntheses in a manner akin to that of photosynthesis.^{9,10}

In contrast to traditional synthetic routes, photoredox and metallaphotoredox catalysts utilize light to drive reactivity under very mild conditions that would otherwise be inaccessible or very costly.¹¹ These methodologies have been mostly employed in the synthesis of small molecules, such as pharmaceuticals and perfumes, but have yet to be translated into mass production of feedstock chemicals or their valorization at large scale.¹² Photocatalytic methods, though highly efficient and mild, proceed via very intricate and meticulously crafted mechanisms that are not easily applicable into broad scope large scale reactions.^{9,10} Translating these technologies to a wider range of chemical manufacturing applications therefore requires new approaches that make the reaction steps more robust and more efficient, as well as the underlying mechanistic knowledge to target these steps. Such efforts are intrinsically highly collaborative as tools and techniques from multiple fields must be utilized. In this context, the Bioinspired Light-Escalated Chemistry (BioLEC) Energy Frontier Research Center was

founded as an interdisciplinary conglomerate of researchers (physicists, physical chemists, biochemists, organic chemists, and engineers) that synergize to advance photoredox catalysis (Figure 1).

This Account summarizes recent collaborative accomplishments in photoredox catalysis through mechanistic understanding, the development of new reactivities, novel applications of new and existing reactions, and the utilization of photoredox catalysis to tackle industrially relevant problems (Figure 1). We first focus on fundamental studies of photocatalysts and intermediates proposed in traditional photoredox catalytic cycles. Then, we discuss the development of new reactions and the optimization of already existing reactions guided by mechanistic studies. Next, we address the development of unique light-harvesting strategies to drive reactivity. Finally, we show how developed methodologies can transform highly available biomass into chemical feedstocks, and how a lightmediated depolymerization mechanism degraded hydroxylated polymers to yield well-defined monomers that can be later functionalized to enter a much sought-after closed-loop recycling cycle without generating stoichiometric waste.

2. SPECTROSCOPIC TOOLS AS PROBES TO INCREASE FUNDAMENTAL UNDERSTANDINGS OF PHOTOREDOX CATALYSIS

Photocatalytic mechanisms are often proposed based on yield analysis (NMR), Stern–Volmer quenching, and electrochemistry. However, these analytical methods generally do not enable a specific step of a complex mechanism to be interrogated, particularly where short-lived reactive intermediates are invoked. Through collaborations, we employed three timeresolved spectroscopies to probe cutting-edge catalytic mechanisms with greater control over species generation and temporal resolution: pulse radiolysis (PR), time-resolved dielectric loss (TRDL), and photoinduced transient absorption (TA) spectroscopy.

2.1. Pulse Radiolysis

The emerging field of metallaphotoredox^{11,13–15} chemistry is exemplified in the Ir/Ni dual photocatalytic system used in cross-coupling reactions to form new C–C and C–heteroatom bonds.^{16–18} The first step in many of these reactions is electron transfer from the Ir(III)* photocatalyst to the Ni(II) precatalyst (e.g., (dtbby)NiBr₂). However, ambiguity remained as to which Ni species is involved in the catalytic cycle, and studying the reduction of (dtbby)NiBr₂ with electrochemistry shows irreversible behavior due to the two-electron reduction followed by subsequent chemistry.¹⁹ Therefore, a different approach was needed. PR is a powerful technique that has yet to be applied in metallaphotoredox catalysis. PR can be thought of as "electrochemistry without the electrode," enabling rapid (nanosecond) one-electron reduction of Ni(II) in solution to unambiguously generate Ni(I) (see Figure 2A).

A collaboration between the MacMillan group and Brookhaven National Laboratory (BNL) generated Ni(I) with PR and then studied its reactivity with TA spectroscopy.²⁰ PR initiates by sending a short pulse (~10 ps) of high energy electrons (9 MeV) through a solution, ionizing the solvent.²¹ Within nanoseconds, charges or excited states can transfer to solutes to create intermediates present in photocatalytic cycles.²² By comparing the Ni(I) spectrum to that obtained with spectroelectrochemistry, we discovered that here, Ni(0) must comproportionate to two Ni(I) molecules. Furthermore,



Figure 2. Application of time-resolved spectroscopies to understand fundamental processes in photoredox catalysis. (A) Pulse radiolysis enables the rapid one-electron reduction of Ni(II) in solution to help elucidate the mechanism of oxidative addition in Ir/Ni dual photocatalytic systems. Adapted from ref 20. Copyright 2021 American Chemical Society. (B) Time-resolved dielectric loss spectroscopy measures the dynamics of the dipole of a common Ir photocatalyst during photoexcitation and finds the counterion to play an important role in excited state reactivity. Adapted with permission from ref 1. Copyright 2022 Springer Nature Limited. (C) Photoinduced transient absorption studies the evolution of the excited state of a Ni photocatalyst that is capable of Ni–Ar bond homolysis from direct excitation. Adapted from ref 32. Copyright 2020 American Chemical Society.

we measured rates of reaction of Ni(I) with a series of electronically distinct aryl iodides. Rates for oxidative addition ranged from 1.3×10^4 to 2.4×10^5 M⁻¹ s⁻¹ depending on the

Hammett value of the para substituent, σ_{para} . Finally, the $[(\text{dtbbpy})\text{NiBr}]_2$ dimer was found to be unreactive with iodobenzene and determined to be mechanistically irrelevant. By generating and characterizing the Ni(I) intermediate, PR helped elucidate this oxidative addition process under synthetically relevant conditions. These studies are of general relevance in Ir/Ni metallaphotoredox catalysis as the formation of elusive Ni(I) species is often postulated as it is also evidenced in the arylation of secondary amines discussed in section 3.

PR is well-suited to such mechanistic studies as it can generate a diverse selection of reactive intermediates such as neutral and charged radicals, H-adducts, triplet excited states, and radical ion pairs. PR can also generate potent redox agents, from the solvated electron to the solvent radical cation, enabling the activation of substrates out of reach to most single-photon photocatalysts. These studies improve fundamental understanding of complex photoredox systems, potentially leading to improved efficiency.

2.2. Time-Resolved Dielectric Loss Spectroscopy

Transition metal complexes are popular for photocatalysis because the formation of their long-lived triplet charge transfer states can drive chemistry. Since many TM complexes are ions, it is likely that the inert counterions can be tuned to enhance photocatalytic reactivity or selectivity. However, the dynamic behavior of counterions following photoexcitation had yet to be studied, as few methods exist to directly characterize ion pairing, let alone response to perturbation.²³ We probed these dynamic counterion effects for the first time and related them to chemical reactivity, providing information about the unique role counterions can play.

TRDL spectroscopy is sensitive to dipoles, whereas other time-resolved spectroscopic methods may probe electronic transitions, bond vibrations, or spin states. Heteroleptic TM complexes exhibit dipole moments in the ground and excited states that can be monitored with TRDL. A collaboration led by the National Renewable Energy Laboratory (NREL) with the Scholes, McCusker, and Knowles groups applied this technique to the popular photocatalyst [Ir[dF(CF₃)ppy]₂(dtbpy)]⁺X⁻ and observed the effect of a weakly (X⁻ = BAr^F₄⁻) versus tightly (X⁻ = PF₆⁻) bound counterion on the dipole and excited-state chemical reactivity.¹

Ir complexes were measured in a microwave cavity (Figure 2B) coupled to a waveguide apparatus which can record the time-dependence of dielectric properties of the solution (real and imaginary). Computational modeling of the cavity response gives the ground state dipole and shows its evolution in the excited state. Dipole dynamics show a sub-nanosecond reorganization of the counterion in response to the dipole change within the complex. Substituting the PF_6^- counterion with the weakly associating $BArF_4^-$ prevented this reorganization and reduced the dipole of the MLCT state. Stern–Volmer quenching studies showed a corresponding decrease in the rates of substrate reduction and increase in the rates of oxidation for PF_6^- relative to $BArF_4^-$.

TRDL is a particularly powerful tool to probe the effect of charge distribution on excited-state reactivity, especially in cases where traditional spectroscopy cannot. We now recognize that the identity of the inert counterion can be optimized when developing a new reaction, particularly in low polarity solvents susceptible to ion pairing effects.

2.3. Photoinduced Transient Absorption Spectroscopy

Ni-photoredox catalysis has also been proposed to proceed via direct photoexcitation^{24–26} or energy transfer based on product analysis.^{27–31} Scholes, Castellano, Doyle, and co-workers investigated a series of photoexcited Ni(^Rbpy)(Ar)Cl derivatives with TA spectroscopy, which can directly generate and probe excited-state dynamics.³² Spectral analysis of Ni(^{t-Bu}bpy)(o-Tol) Cl (Figure 2C) revealed an initial MLCT state evolves into a ³dd excited state over the first 15 ps (τ = 5.5 ps). The excited Ni(II) complexes then undergo Ar-Ni bond homolysis to generate aryl radicals and Ni(I), which offers a novel mechanism to initiate catalysis by Ni(I) in the absence of an exogenous photocatalyst.³³ TA spectroscopy provided vital information that will enable this new mode of reactivity to be exploited. Similar spectroscopic methods have been applied to Ni-photoredox to differentiate energy versus electron transfer in Ni-mediated C-O coupling²⁹ and dark Ni(I)/Ni(III) cycles in C-N crosscoupling reactions.³⁴ Overall, we demonstrated that collaborative efforts using unique spectroscopic methodologies provide valuable insight into, and potentially inform future, photocatalytic reactions. More sophisticated techniques, such as multidimensional spectroscopy, could potentially further disentangle the ultrafast mechanisms behind the initial steps of photoredox catalysis.

3. DEVELOPMENT OF NEW REACTIVITIES GUIDED BY MECHANISTIC INSIGHTS

Gaining mechanistic insight into the elementary steps of photoredox catalysis facilitates the advancement and development of new and more efficient reactions in organic synthesis which translates to other fields. The spectroscopic techniques discussed in the previous section enable in-depth mechanistic investigations of elementary steps in photocatalyzed reactions. These mechanistic understandings can be leveraged and translated into the development of new reactivities or an improvement of existing ones through interdisciplinary collaborative efforts.

A notable example of how mechanistic understandings can lead to the improvement of existing reactions is the design of a bioinspired Ir complex with a proton-coupled electron transfer (PCET)-based ligand for the reductive cleavage of *N*-(acyloxy)phthalimide redox-active esters (Figure 3A).² These efforts were carried out by the Knowles, Moore, Scholes, and Rumbles groups in 2021. A number of photocatalytic transformations suffer from limited scopes and/or quantum efficiencies due to charge recombination (CR) issues between the oxidized/ reduced photocatalyst and the radical species being formed.³⁵ In order to increase reaction efficiencies, the rate of the forward reaction must be such that it is competitive with the rate of bimolecular CR processes. Inspired by electron relays present in Photosystem II that can transfer oxidizing/reducing equivalents as a means to suppress CR events,³⁶ we designed a ligand that contains a BIP (benzimidazole-phenol) moiety that is able to, upon excitation of the photocatalyst, undergo a PCET process to populate an intramolecular charge separated state (CSS) that inhibits the recombination of the subsequent intermolecular CSS (Figure 3B). Utilizing the methyl viologen dication as a substrate surrogate, we were able to demonstrate that the CR process is, in fact, reduced by 24-fold when compared to the parent photocatalyst, $[Ir(dF(CF_3)-ppy)_2(bpy)](PF_6)$. Excitingly, the CR reduction led to a 2-fold increase in quantum



Figure 3. PCET-based ligand to address intramolecular charge recombination. (A) General reaction and mechanism for the reduction of redox-active esters. (B) Ligand design employed to hinder intermolecular charge recombination and comparison of reaction efficiencies of the traditional $[Ir(dF(CF_3)-ppy)_2(bpy)](PF_6)$ photocatalyst against $[Ir(dF(CF_3)-ppy)_2(BIP-bpy)](PF_6)$. Adapted from ref 2. Copyright 2021 American Chemical Society.

yield efficiency for the photocatalytic reduction of a *N*-(acyloxy)phthalimide redox-active ester.

In 2020, the MacMillan and Scholes groups collaborated to investigate the mechanism and improve upon a metallaphotocatalyzed C–N cross-coupling reaction (Figure 4A).³⁴ First, Stern–Volmer luminescent quenching experiments of the standard Ir photocatlyst led to the assignment of DABCO as the primary quencher of its excited state. In combination with nanosecond–microsecond pump–probe spectroscopies, alongside with spectroelectrochemistry, we were able to show that, upon quenching, the formation of Ir(II) and DABCO^{•+} species would follow. Moreover, these species were determined to be pertinent as the formation of Ir(II) was shown to reduce Ni(II)



Figure 4. Photocatalyst redesign strategies for C–N cross-coupling. (A) General reaction and mechanism of a C–N coupling reaction. (B) Photocatalyst redesign, Stern–Volmer plot for DABCO quenching of the photocatalyst, and rates of product formation of a series of Ir photocatalyst as a function of their redox potentials. Adapted from ref 34. Copyright 2020 American Chemical Society.

to a catalytically relevant and elusive Ni(I) species (as demonstrated in the previous section) that can undergo oxidative addition. Additionally, the efficiency of this step was determined to be crucial for the initiation and perpetuation of the Ni(I)/Ni(III) cycle. Hence, new Ir(III) photocatalysts of increasing electron density on the π^* -localized bipyridine ligands were synthesized and characterized to increase the reductive capability of the intermediate Ir(II) complexes. In particular, the more highly reducing photocatalysts developed in the series led to a 30-fold rate increase and a 10-fold increase in quantum yield. Furthermore, the redesign of the Ir photocatalyst resulted in an expanded scope that includes synthetically useful yields of product formation with challenging drug-like aryl halide substrates previously inaccessible. Overall, the merger of

these techniques has demonstrated the chemical advancements that come about from the understanding of the intricacies of the underlying mechanisms of the reactions.

Doyle and co-workers reported a dual phosphine and Ir photocatalytic pathway to construct $C(sp^3)-N$ bonds via an intermolecular anti-Markovnikov hydroamination of unactivated olefins with primary sulfonamides (Figure 5A).³⁷



these experiments are inconsistent with a PCET pathway supporting a phorphoranyl darical mechanism

Figure 5. α -Scission of phosphoryl radicals as precursors of N-centered radicals for olefin hydroamination. (A) General reaction and key mechanistic steps. (B) Stern–Volmer experiments consistent with phosphoranyl radical formation. Adapted from ref 37. Copyright 2021 American Chemical Society.

Mechanistic investigations support that an excited photocatalyst oxidizes the phosphine to the corresponding radical cation and forms a phosphoranyl radical intermediate upon nucleophilic addition of the sulfonamide which subsequently generates the sulfonamidyl N-centered radical via α -scission of the P–N bond. This work evaluates multiple mechanistic hypotheses of the transformation and systematically addresses the support and lack thereof for each pathway. One alternative hypothesis considered for the sulfonamide N–H activation was concerted PCET. The results of Stern–Volmer experiments were inconsistent with a PCET mechanism and supported the proposed pathway (Figure 5B). A better understanding of the reactivity was gained by using spectroscopic experiments, and this cocatalytic system is under active investigation for the activation of other challenging substrates.

The examples presented in this section show how elucidating the elementary steps of a mechanism can provide valuable clues for reaction optimizations as well as for the development of reactions that were previously inaccessible. In particular, photocatalyst design can have a tremendous impact on reaction outcomes. Understanding the limitations of the elementary steps of these complex mechanisms can provide clues for the design of more efficient photocatalysts that can increase reaction yields, rates, and even expand the scope to more challenging substrates. In particular, the design of more efficient photocatalysts, as well as the implementation of spectroscopic tools to provide insight into the elementary steps of a transformation, may lead to the development of, for example, more efficient depolymerization strategies such as the ones described in section S.

4. LOW-ENERGY LIGHT-HARVESTING STRATEGIES FOR CATALYTIC REACTIVITY

Light absorption is the key step needed for photocatalysis to occur.⁹ Many catalysts suffer from light-limited operation due to their small absorption bandwidths and low molar absorptivities and are furthermore susceptible to photodamage and nonspecific excitation.^{10,38} Previous advancements in widening the absorption bandwidth utilized such tactics as conjugated transition-metal complexes,^{39,40} upconverted light to excite a catalyst,⁴¹⁻⁴³ and direct excitation of the triplet state of a catalyst.^{44,45} Particularly, expanding a catalyst's available absorption window to lower energies can avoid nonspecific excitation of optical filters such as substrates and cocatalysts, improve a catalyst's implementation in relevant reactions such as biological cross-linking in vivo, reduce potential damage from irradiation with high-energy light, and improve the scalability of reactions due to increased penetration depths compared to high energy light.^{38,46} We designed, investigated, and demonstrated two unique light-harvesting strategies: energy transfer from a dedicated light-harvesting protein to a tethered Ru photocatalyst and triplet-triplet annihilation for direct initiation of freeradical polymerization, in the context of photocatalytic reactions (Figure 6A and C).^{3,47}

Schlau-Cohen, MacMillan, Doyle, Castellano, and co-workers utilized a low-energy absorbing photosynthetic light-harvesting protein to excite a photocatalyst through energy transfer and catalyze two radical initiation reactions.³ Natural organisms partition the light-harvesting and reactive processes of photosynthesis into distinct structural components.⁴⁸ This modulation occurs because these processes are contradictory, one requiring harnessing and storing vast amounts of energy and the other quickly dispersing energy toward chemical reactivity. The authors sought to apply this design principle to the development of a photocatalytic bioconjugated system in the context of photoredox catalysis. Due to the intrinsic interdisciplinary nature of the work proposed, a collaboration between several groups was necessary to make the developments required to implement this chemistry. The Schlau-Cohen group conjugated the light-harvesting protein R-phycoerythrin (RPE) to the photocatalyst [Ru(bpy)₃]²⁺ to develop the biohybrid catalyst $RPE-(Ru)_n$ and examined the energy transfer time scale between the components using fluorescence lifetime measurements, transient absorption spectroscopy, and Förster theory (Figure 6A). The Castellano group performed nanosecond transient absorption measurements, which provided concrete evidence that RPE could transfer energy to $[Ru(bpy)_3]^{2+}$; excitation at wavelengths at which only RPE absorbs resulted in spectral



Figure 6. Strategies for advantageous light absorption. (A) Energy transfer from a conjugated broadly absorbing light-harvesting protein enables lowenergy light to excite a reactive catalyst. (B) Spectral features of both the light-harvesting protein and catalyst were observed through transient absorption spectroscopy when exciting only the protein, evidencing energy transfer. The biohybrid enabled or enhanced product yields in a photocatalytic thiol-ene test reaction after 2 h. ^a12 h reaction time. Adapted with permission from ref 3. Copyright 2022 Elsevier. (C) Triplet-triplet annihilation enables low-energy light to excite and initiate polymerization between methyl acrylate and trimethylolpropane triacrylate. (D) Transient absorption spectroscopy and Stern–Volmer quenching experiments lend evidence toward triplet-triplet annihilation as the operative mechanism. Adapted with permission from ref 47. Copyright 2020 Elsevier.

signatures of $[Ru(bpy)_3]^{2+}$, indicating energy transfer had occurred (Figure 6B). Finally, the MacMillan and Doyle groups demonstrated the biohybrid for synthetic competency through photocatalytic thiol—ene coupling and cysteinyl desulfurization test reactions. Here, reactivity was enhanced at green wavelengths of common absorption for the components and enabled at red wavelengths where the catalyst alone does not absorb (Figure 6B). This strategy demonstrated the viability of uphill energy transfer from a conjugated light-harvesting structure as a means of enabling the low-energy excitation of a photocatalyst. Similar to the annihilation work described below, these biohybrids offer increased absorption bandwidths, noncompetitive absorption with substrates and cocatalysts, and applications in biological systems.^{46,49}

Castellano and co-workers utilized triplet—triplet annihilation to directly catalyze a free-radical polymerization of methyl acrylate (MA) or trimethylolpropane triacrylate (TMPTA) with low-energy light.⁴⁷ Triplet—triplet annihilation is a multiphoton excitation mechanism which combines the exciting power of two low-energy photons and allows high-energy light to be emitted or high-energy chemistry to be performed. The authors excited the lower singlet excited states (S_1) of two ZnTPP (zinc(II) meso-tetraphenylporphyrin) molecules, which rapidly intersystem crossed to their respective triplet states (T_1) (Figure 6C). Two of these triplet excited photoinitiators then annihilated to form a ground state (S_0) molecule and a high-energy excited singlet state (S_2) molecule that directly initiates the free-radical polymerization of TMPTA or MA. Using transient absorption spectroscopy and time-resolved and steady-state fluorescence measurements, the authors determined that only the T₁ states and the S_2 state were mechanistically relevant (Figure 6D), as their spectral features were quenched in the presence of TMPTA, lending evidence to triplet-triplet annihilation as the operative mechanism. The S2 state of ZnTPP directly catalyzes the radical polymerization of TMPTA through electron transfer. The polymerization was monitored by witnessing the decay of the C=C IR band of the monomer over time, and EPR spectroscopic studies supported a mechanism proceeding via radical-initiated upconverted light. This strategy demonstrated the viability of utilizing upconverted

light to directly catalyze reactions, offering excitation with lowenergy sources, fewer competing side reactions, and less complexity as the upconverter simultaneously acts as the photoinitiator.

As part of our overarching goal to improve photocatalysis, we aim to expand the possibilities for innovative light absorption and confirm viability of developed strategies. By applying triplet-triplet annihilation and dedicated light-harvesting machinery to photocatalytic processes, we demonstrated that low-energy visible light can be used for irradiation and that the possibility of reactions occurring in aqueous and biological environments can one day be realized. In collaboration with the synthetic organic groups, we are currently exploring the generalizability of these strategies including conjugating lightharvesters to the chemically remarkable photoenzymatic mutant, Gluconobacter Ene-Reductase T36A, developed by Hyster and co-workers.⁵⁰ We hope to expand these methods to further reactions and systems, such as developing biohybrids containing earth-abundant metal-centered catalysts, expanding the number of reactions which can be directly catalyzed by an upconverting system, and exploiting the potential power of multiphoton excitation and multielectron reactivity.⁵

5. DEPOLYMERIZATION STRATEGIES ENABLED BY PHOTOREDOX CATALYSIS VIA PCET

5.1. Depolymerization of Native Lignin

Due to increasing energy demand and its impact on Earth's climate, moving away from nonrenewable energy sources, such as fossil fuels, is critical. Biomass is an appealing alternative, as it is considered the most abundant renewable source of carbon, biofuels, and chemical feedstocks (Figure 7A).¹² In particular, lignin, a carbon-neutral nonedible polymer, is generated in large quantities by agricultural activity worldwide.⁵² The depolymerization of this material into useful monomeric products, however, remains challenging due to the high temperatures and harsh conditions required to break the C–C and C–O linkages that hold the biopolymer together. Light-driven depolymerization of this biomass source is therefore an attractive alternative as it can be achieved under mild conditions without generating stoichiometric waste.⁵³

Knowles and co-workers employed photoredox catalysis to depolymerize native lignin via a proton-coupled electron transfer (PCET) activation step.⁵⁴ The mechanism for the degradation of this polymer starts with excitation of the Ir(III) photocatalyst into a highly oxidizing Ir(III)* excited state (Figure 7B). In the presence of a base, the hydroxy groups in the polymer can quench this excited state via a PCET step, generating an Ir(II) species and an alkoxy radical.55,56 This unstable radical undergoes C-C bond β -scission to yield an aldehyde and an alkyl radical that can, through a hydrogen atom transfer (HAT) step, yield depolymerized product. The HAT reagent is regenerated by reoxidation of the Ir(II) species followed by a proton transfer step closing the catalytic cycle. This one-step, redox neutral reaction proceeds at room temperature without the generation of stoichiometric waste. This approach yielded monomers with efficiencies that range from 57 to 98% for model lignin dimeric systems that display β -O-4 linkages. However, the cleavage of plant-derived native lignin is significantly more challenging than these model systems and small molecule lignin fractions, meaning literature examples are scarcely found. Of note, Stephenson and co-workers developed a two-step depolymerization method that consists



Figure 7. Depolymerization of biomass. (A) Proposed mechanism for the depolymerization of lignin. (B) Top: biomass as a source of chemical feedstock through solar light driven oxidation.⁵³ Bottom: schematic representation of native lignin and the products obtained through photocatalytic depolymerization via PCET. Adapted from ref 54. Copyright 2020 American Chemical Society.

of the electrocatalytic oxidation of native lignin followed by photocatalytic cleavage promoted by an Ir phocatalyst that

н

pubs.acs.org/accounts

afforded 2.4 wt % of monomeric material through C–O cleavage of the β -O-4 linkages.⁵⁷ In comparison, the photoredox C–C cleavage via PCET method developed by the Knowles group yielded 4 wt % of monomeric material through the preparative scale depolymerization of ponderosa lignin in a manner complementary to Stephenson's two-step approach. Excitingly, native grand fir lignin degraded via PCET was estimated to undergo a minimum of 41% bond cleavage of all β -aryl ether bonds present through derivatization followed by reductive cleavage analysis.

5.2. Depolymerization of Hydroxylated Polymers

The above-described method represents a very promising strategy that could be adapted for the depolymerization of other materials of great societal interest such as plastics. Even though the development and widespread usage of synthetic polymers has had an unarguably positive influence in societal development, the accompanying environmental effects still represent a challenge that needs to be addressed.⁵⁸ The recycling strategies of many plastics require multiple C-C bond-breaking steps that are typically hard to access and degrade, often yielding complex mixtures that cannot easily be repolymerized. Hydroxylated polymers, however, are a subclass of plastics that display hydroxyl groups in the backbone which renders them susceptible to depolymerization strategies that proceed via the formation of an alkoxy radical and subsequent C-C bond breaking step. The Knowles group sought to depolymerize these types of polymers via a PCET enabled photoredox catalytic cycle.⁴ The degradation of a phenoxy resin yielded the desired monomers in up to 63% yield (Figure 8A). This polymeric resin is typically found in mixtures, and excitingly, degradation of this resin in the presence of added commodity plastics such as poly(vinyl chloride) ($M_{\rm w} \sim 233\ 000\ {\rm g/mol}$), polystyrene ($M_{\rm w} \sim$ 192 000 g/mol), and poly(methyl methacrylate) ($M_w \sim 120\ 000$ g/mol) did not significantly affect the yields of monomer obtained (50-54%) as evidenced by GC analysis. Interestingly, the monomeric product can be easily converted into bisphenol A with a 93% yield, which is a precursor for many other commercial polymers. Gratifyingly, this methodology can be extended to the degradation of hydroxylated homo and copolymers with *p*-methoxyphenyl (PMP) and silylether (OTBS) substituents at the β -position due to stabilization of the C-centered radical formed upon β -scission. Moreover, the degradation products can be functionalized to afford bifunctional monomers as chemical feedstocks creating a closed-loop of polymerization/depolymerization of recyclable polymers (Figure 8B).

These examples show how photoredox catalysis provides a mild alternative to traditional polymer degradation strategies for the production of chemical feedstocks from abundant biomass as well as creates a closed recycling loop for synthetic polymers containing hydroxyl groups. Implementation of PR and ultrafast spectroscopies described in section 2 may provide further mechanistic insights into the elementary steps involved in the degradation of polymers. Additionally, the development of new activation modes for depolymerization reactions, such as the multi-photo processes described in section 3, might allow us to expand the scope to the degradation of more challenging nonrecyclable plastics.

6. CONCLUSIONS

Light-driven chemistry has the potential to revolutionize the chemical industry as it provides alternative reactivity pathways



Figure 8. Depolymerization of hydroxylated polymers. (A) Proposed degradation/repolymerization cycle for the recycling of hydroxylated polymers via product functionalization and photocatalytic depolymerization. (B) Monomeric products obtained by photocatalytic depolymerization via PCET of phenoxy resins and hydroxylated copolymers and their product distributions. Adapted with permission from ref 4. Copyright 2021 American Chemical Society.

that lead to the formation of products that are either unattainable through traditional routes or energetically costly due to the high temperatures and pressures required to obtain them. The advancement of this field, however, requires collaborative efforts in order to grow due to the intrinsically interdisciplinary nature of the work required to understand and exploit all aspects of photoredox chemistry. The use of techniques such as PR, TRDL, and TA spectroscopies are not routinely employed by organic chemists, but collaborations across the BioLEC team allowed us to investigate elementary steps in photoredox catalysis that would otherwise remain obscure. These results led us to discover new reactivities and new optimization variables (i.e., the anion of the photocatalyst modulates ES dynamics) of importance to synthetic chemists. Additionally, we developed new photocatalysts that improve the efficiency of C-N coupling reactions guided by spectroscopic techniques as well as C-N cross coupling reactions that proceed through the formation of N-centered radicals generated via a α scission of a phosphoranyl radical intermediate. Moreover, we successfully employed two light harvesting strategies, triplettriplet annihilation and bioconjugation, to circumvent nonspecific absorption issues in the contexts of polymerization and thiol-ene coupling reactions with enhanced reactivity. These could have a tremendous impact on scaled-up industrial applications as light penetration increases for these low-energy light-induced processes facilitating chemistry in batch reactors. Furthermore, the discovery of the activation of alcohols through a PCET mechanism that results in C–C β -scission of an alkoxy radical led us to the development of a highly efficient method for the challenging depolymerization of native lignin and the depolymerization of industrially relevant hydroxylated polymers. Excitingly, the products of the latter lead to the formation of feedstock chemicals which we leveraged to establish a closedrecycling loop through post-functionalization. These highlighted collaborative efforts developed by BioLEC opened new avenues of exploration in the context of photoredox catalysis that have advanced the field in unprecedented directions.

AUTHOR INFORMATION

Corresponding Authors

- David W. C. MacMillan Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States; orcid.org/0000-0001-6447-0587; Email: dmacmill@ princeton.edu
- Gregory D. Scholes Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States; orcid.org/0000-0003-3336-7960; Email: gscholes@ princeton.edu

Authors

- Agustin Millet Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States
- **Paul T. Cesana** Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States
- Kassandra Sedillo Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States
- Matthew J. Bird Chemistry Division, Brookhaven National Laboratory, Upton, New York 11973, United States; orcid.org/0000-0002-6819-5380
- Gabriela S. Schlau-Cohen Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States; orcid.org/0000-0001-7746-2981
- Abigail G. Doyle Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States; Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095, United States;
 orcid.org/0000-0002-6641-0833

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.accounts.2c00083

Notes

The authors declare no competing financial interest.

Biographies

Agustin Millet is currently a Distinguished Postdoctoral Fellow of the BioLEC EFRC. He received his B.S. in chemistry from the University of Buenos Aires, Argentina after which he obtained his Ph.D. in 2019 under the supervision of Prof. Kim R. Dunbar at Texas A&M University. In the laboratory of Prof. Dave W. C. MacMillan, his current research focuses on the development of new metallaphotoredox methodologies.

Paul T. Cesana received his B.S. in chemistry from the University of Rhode Island in 2019 and is now a Ph.D. student in Prof. Gabriela S. Schlau-Cohen's laboratory at the Massachusetts Institute of Technology (MIT). His research focuses on developing strategies to improve light-harvesting by photocatalysts inspired by photosynthetic organisms.

Kassandra Sedillo received her B.S. in chemistry from Colorado State University in 2015 and conducted research in the lab of Prof. Tom Rovis. Afterward, she worked as a medicinal chemist at Gilead Sciences in Seattle, Washington and is now a Ph.D. candidate at Princeton University coadvised by Prof. Abigail G. Doyle (currently at UCLA) and Prof. Robert R. Knowles. Her current research focuses on methodology development for direct bond activation using phosphine and photoredox catalysis.

Matthew J. Bird is an Associate Chemist at Brookhaven National Laboratory. He received his Ph.D. under the direction of Prof. Henning Sirringhaus at the University of Cambridge in 2011. His research focuses on the use of pulse radiolysis to study electron and energy processes relevant to solar energy conversion as well as fundamental studies of radiation and electrochemistry.

Gabriela S. Schlau-Cohen is an Associate Professor of Chemistry at the Massachusetts Institute of Technology (MIT) in Cambridge, MA. She received her Ph.D. under the direction of Prof. Graham R. Fleming at the University of California, Berkeley in 2011 and completed postdoctoral research at Stanford University under Prof. W. E. Moerner in 2014. Her research focuses on the spectroscopic investigation of photosynthetic systems and the implementation of this knowledge to new technologies and applications.

Abigail G. Doyle is currently the Saul Winstein Chair in Organic Chemistry at the University of California, Los Angeles. She received her Ph.D. from Harvard University under the direction of Prof. Eric Jacobsen and served as a Professor of Chemistry at Princeton University from the start of her independent career in 2008 until 2021. Her research focuses on contributing new approaches to chemical synthesis and catalysis, especially using Ni-centered species.

David W. C. MacMillan is the James S. McDonnell Distinguished University Chair of chemistry at Princeton University. He received his Ph.D. under Prof. Larry Overman at the University of California, Irvine in 1996 and was a postdoctoral scholar with Prof. David Evans at Harvard University. In 2021, he was awarded the Nobel Prize in Chemistry with Prof. Benjamin List for the development of the field of asymmetric organocatalysis. His research focuses on wide range of organic chemistry transformations that includes the development of new methodologies enabled by photoredox and metallaphotoredox catalysis.

Gregory D. Scholes is currently the William S. Tod Professor of Chemistry at Princeton University, the chair of Princeton's chemistry department, editor-in-chief of the Journal of Physical Chemistry Letters, and director of the BioLEC EFRC. He received his Ph.D. from the University of Melbourne in 1994 under the direction of Prof. Ken Ghiggino and was a postdoctoral scholar at the University of California, Berkeley under Prof. Graham Fleming. His research spectroscopically investigates natural and bioinspired photosynthetic systems for new applications in light harvesting and catalysis.

REFERENCES

(1) Earley, J. D.; Zieleniewska, A.; Ripberger, H. H.; Shin, N. Y.; Lazorski, M. S.; Mast, Z. J.; Sayre, H.; Knowles, R. R.; McCusker, J. K.; Scholes, G. D.; Reid, O. G.; Rumbles, G. Ion Pair Reorganization Regulates Reactivity in Photoredox Catalysts.1. *Nat. Chem.* **2022**, DOI: 10.1038/s41557-022-00911-6.

(2) Sayre, H.; Ripberger, H. H.; Odella, E.; Zieleniewska, A.; Heredia, D. A.; Rumbles, G.; Scholes, G. D.; Moore, T. A.; Moore, A. L.; Knowles, R. R. PCET-Based Ligand Limits Charge Recombination with an Ir(III) Photoredox Catalyst. J. Am. Chem. Soc. 2021, 143, 13034–13043.

(3) Cesana, P. T.; Li, B. X.; Shepard, S. G.; Ting, S. I.; Hart, S. M.; Olson, C. M.; Martinez Alvarado, J. I.; Son, M.; Steiman, T. J.; Castellano, F. N.; Doyle, A. G.; MacMillan, D. W. C.; Schlau-Cohen, G. S. A Biohybrid Strategy for Enabling Photoredox Catalysis with Low-Energy Light. *Chem* **2022**, *8*, 174.

(4) Nguyen, S. T.; McLoughlin, E. A.; Cox, J. H.; Fors, B. P.; Knowles, R. R. Depolymerization of Hydroxylated Polymers via Light-Driven C-C Bond Cleavage. J. Am. Chem. Soc. **2021**, 143, 12268–12277.

(5) *Guide to the Business of Chemistry*; American Chemistry Council: Washington DC, 2019.

(6) Sayre, H. J.; Tian, L.; Son, M.; Hart, S. M.; Liu, X.; Arias-Rotondo, D. M.; Rand, B. P.; Schlau-Cohen, G. S.; Scholes, G. D. Solar Fuels and Feedstocks: The Quest for Renewable Black Gold. *Energy Environ. Sci.* **2021**, *14*, 1402–1419.

(7) Harrington, R. This Incredible Fact Should Get You Psyched About Solar Power. *Business Insider*, September 29. 2015.

(8) National Renewable Energy Laboratory. Best Research-Cell Efficiency Chart; NREL. https://www.nrel.gov/pv/cell-efficiency.html.

(9) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. J. Org. Chem. 2016, 81, 6898-6926.

(10) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322–5363.

(11) Chan, A. Y.; Perry, I. B.; Bissonnette, N. B.; Buksh, B. F.; Edwards, G. A.; Frye, L. I.; Garry, O. L.; Lavagnino, M. N.; Li, B. X.; Liang, Y.; Mao, E.; Millet, A.; Oakley, J. V.; Reed, N. L.; Sakai, H. A.; Seath, C. P.; MacMillan, D. W. C. Metallaphotoredox: The Merger of Photoredox and Transition Metal Catalysis. *Chem. Rev.* **2022**, *122*, 1485–1542.

(12) Tuck, C. O.; et al. Valorization of Biomass: Deriving More Value from Waste. *Science*. **2012**, 337, 695–699.

(13) Twilton, J.; Le, C. C.; Zhang, P.; Shaw, M. H.; Evans, R. W.; MacMillan, D. W. C. The Merger of Transition Metal and Photocatalysis. *Nat. Rev. Chem.* **2017**, *1*, 0052.

(14) Levin, M. D.; Kim, S.; Toste, F. D. Photoredox Catalysis Unlocks Single-Electron Elementary Steps in Transition Metal Catalyzed Cross-Coupling. *ACS Cent. Sci.* **2016**, *2*, 293–301.

(15) Zhu, C.; Yue, H.; Chu, L.; Rueping, M. Recent Advances in Photoredox and Nickel Dual-Catalyzed Cascade Reactions: Pushing the Boundaries of Complexity. *Chem. Sci.* **2020**, *11*, 4051–4064.

(16) Zuo, Z.; Ahneman, D. T.; Chu, L.; Terrett, J. A.; Doyle, A. G.; MacMillan, D. W. C. Merging Photoredox with Nickel Catalysis: Coupling of α -Carboxyl Sp3-Carbons with Aryl Halides. *Science*. **2014**, 345, 437–440.

(17) Tellis, J. C.; Primer, D. N.; Molander, G. A. Single-Electron Transmetalation in Organoboron Cross-Coupling by Photoredox/ Nickel Dual Catalysis. *Science.* **2014**, *345*, 433–436.

(18) Corcoran, E. B.; Pirnot, M. T.; Lin, S.; Dreher, S. D.; Dirocco, D. A.; Davies, I. W.; Buchwald, S. L.; Macmillan, D. W. C. Aryl Amination Using Ligand-Free Ni(II) Salts and Photoredox Catalysis. *Science.* **2016**, 353, 279–283.

(19) Kawamata, Y.; Vantourout, J. C.; Hickey, D. P.; Bai, P.; Chen, L.; Hou, Q.; Qiao, W.; Barman, K.; Edwards, M. A.; Garrido-Castro, A. F.; Degruyter, J. N.; Nakamura, H.; Knouse, K.; Qin, C.; Clay, K. J.; Bao, D.; Li, C.; Starr, J. T.; Garcia-Irizarry, C.; Sach, N.; White, H. S.; Neurock, M.; Minteer, S. D.; Baran, P. S. Electrochemically Driven, Ni-Catalyzed Aryl Amination: Scope, Mechanism, and Applications. *J. Am. Chem. Soc.* **2019**, *141*, 6392–6402.

(20) Till, N. A.; Oh, S.; MacMillan, D. W. C.; Bird, M. J. The Application of Pulse Radiolysis to the Study of Ni(I) Intermediates in Ni-Catalyzed Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2021**, *143* (25), 9332–9337.

(21) Wishart, J. F.; Cook, A. R.; Miller, J. R. The LEAF Picosecond Pulse Radiolysis Facility at Brookhaven National Laboratory. *Rev. Sci. Instrum.* **2004**, *75*, 4359–4366.

(22) Grills, D. C.; Polyansky, D. E.; Fujita, E. Application of Pulse Radiolysis to Mechanistic Investigations of Catalysis Relevant to Artificial Photosynthesis. *ChemSusChem.* **2017**, *10*, 4359–4373.

(23) Marcus, Y.; Hefter, G. Ion Pairing. *Chem. Rev.* **2006**, *106*, 4585–4621.

(24) Lim, C.-H.; Kudisch, M.; Liu, B.; Miyake, G. M. C-N Cross-Coupling via Photoexcitation of Nickel–Amine Complexes. J. Am. Chem. Soc. 2018, 140, 7667–7673.

(25) Shen, X.; Li, Y.; Wen, Z.; Cao, S.; Hou, X.; Gong, L. A Chiral Nickel DBFOX Complex as a Bifunctional Catalyst for Visible-Light-Promoted Asymmetric Photoredox Reactions. *Chem. Sci.* **2018**, *9*, 4562–4568.

(26) Shields, B. J.; Kudisch, B.; Scholes, G. D.; Doyle, A. G. Long-Lived Charge-Transfer States of Nickel(II) Aryl Halide Complexes Facilitate Bimolecular Photoinduced Electron Transfer. *J. Am. Chem. Soc.* **2018**, *140*, 3035–3039.

(27) Heitz, D. R.; Tellis, J. C.; Molander, G. A. Photochemical Nickel-Catalyzed C–H Arylation: Synthetic Scope and Mechanistic Investigations. J. Am. Chem. Soc. **2016**, 138, 12715–12718.

(28) Kudisch, M.; Lim, C.-H.; Thordarson, P.; Miyake, G. M. Energy Transfer to Ni-Amine Complexes in Dual Catalytic, Light-Driven C–N Cross-Coupling Reactions. J. Am. Chem. Soc. **2019**, *141*, 19479–19486.

(29) Tian, L.; Till, N. A.; Kudisch, B.; MacMillan, D. W. C.; Scholes,
G. D. Transient Absorption Spectroscopy Offers Mechanistic Insights for an Iridium/Nickel-Catalyzed C-O Coupling. J. Am. Chem. Soc. 2020, 142, 4555-4559.

(30) Ricci, A. Modern Amination Methods; Wiley-VCH: Weinheim, 2000.

(31) Welin, E. R.; Le, C.; Arias-Rotondo, D. M.; McCusker, J. K.; MacMillan, D. W. C. Photosensitized, Energy Transfer-Mediated Organometallic Catalysis through Electronically Excited Nickel(II). *Science.* 2017, 355, 380–385.

(32) Ting, S. I.; Garakyaraghi, S.; Taliaferro, C. M.; Shields, B. J.; Scholes, G. D.; Castellano, F. N.; Doyle, A. G. ³d-d Excited States of Ni(II) Complexes Relevant to Photoredox Catalysis: Spectroscopic Identification and Mechanistic Implications. *J. Am. Chem. Soc.* **2020**, *142*, 5800–5810.

(33) Cagan, D. A.; Stroscio, G. D.; Cusumano, A. Q.; Hadt, R. G. Multireference Description of Nickel–Aryl Homolytic Bond Dissociation Processes in Photoredox Catalysis. *J. Phys. Chem. A* **2020**, *124*, 9915–9922.

(34) Till, N. A.; Tian, L.; Dong, Z.; Scholes, G. D.; MacMillan, D. W. C. Mechanistic Analysis of Metallaphotoredox C–N Coupling: Photocatalysis Initiates and Perpetuates Ni(I)/Ni(III) Coupling Activity. J. Am. Chem. Soc. **2020**, 142, 15830–15841.

(35) Gould, I. R.; Farid, S. Dynamics of Bimolecular Photoinduced Electron-Transfer Reactions. *Acc. Chem. Res.* **1996**, *29*, 522–528.

(36) Blankenship, R. E. Molecular Mechanisms of Photosynthesis; Blackwell Science: London, 2002.

(37) Chinn, A. J.; Sedillo, K.; Doyle, A. G. Phosphine/Photoredox Catalyzed Anti-Markovnikov Hydroamination of Olefins with Primary Sulfonamides via α -Scission from Phosphoranyl Radicals. *J. Am. Chem. Soc.* **2021**, *143*, 18331–18338.

(38) Zhu, S.; Wang, D. Photocatalysis: Basic Principles, Diverse Forms of Implementations and Emerging Scientific Opportunities. *Adv. Energy Mater.* **2017**, *7*, 1700841.

(39) Yamazaki, Y.; Rohacova, J.; Koike, K.; Ishitani, O. Synthesis and Light-Harvesting Functions of Ring-Shaped Re(I) Trinuclear Complexes Connected with an Emissive Ru(II) Complex. *JACS Au* **2021**, *1*, 294–307.

(40) Yamazaki, Y.; Ishitani, O. Synthesis of Os(II)-Re(i)-Ru(II) Hetero-Trinuclear Complexes and Their Photophysical Properties and Photocatalytic Abilities. *Chem. Sci.* **2018**, *9*, 1031–1041.

(41) Ravetz, B. D.; Pun, A. B.; Churchill, M.; Congreve, D. N.; Rovis, T.; Campos, L. M. Photoredox Catalysis Using Infrared Light via Triplet Fusion Upconversion. *Nature* **2019**, *565*, 343–346.

(42) Glaser, F.; Kerzig, C.; Wenger, O. S. Sensitization-Initiated Electron Transferviaupconversion: Mechanism and Photocatalytic Applications. *Chem. Sci.* **2021**, *12*, 9922–9933.

(43) Bilger, J. B.; Kerzig, C.; Larsen, C. B.; Wenger, O. S. A Photorobust Mo(0) Complex Mimicking $[Os(2,2'-Bipyridine)_3]^{2+}$ and Its Application in Red-to-Blue Upconversion. *J. Am. Chem. Soc.* **2021**, 143, 1651–1663.

(44) Reischauer, S.; Pieber, B. Emerging Concepts in Photocatalytic Organic Synthesis. *iScience* **2021**, *24*, 102209.

(45) Sellet, N.; Cormier, M.; Goddard, J.-P. The Dark Side of Photocatalysis: Near-Infrared Photoredox Catalysis for Organic Synthesis. *Org. Chem. Front.* **2021**, *8*, 6783–6790.

(46) Ryu, K. A.; Kaszuba, C. M.; Bissonnette, N. B.; Oslund, R. C.; Fadeyi, O. O. Interrogating Biological Systems Using Visible-Light-Powered Catalysis. *Nat. Rev. Chem.* **2021**, *5*, 322–337.

(47) Awwad, N.; Bui, A. T.; Danilov, E. O.; Castellano, F. N. Visible-Light-Initiated Free-Radical Polymerization by Homomolecular Triplet-Triplet Annihilation. *Chem.* **2020**, *6*, 3071–3085.

(48) Mirkovic, T.; Ostroumov, E. E.; Anna, J. M.; van Grondelle, R.; Govindjee; Scholes, G. D. Light Absorption and Energy Transfer in the Antenna Complexes of Photosynthetic Organisms. *Chem. Rev.* 2017, 117, 249–293.

(49) Liu, W.; Watson, E. E.; Winssinger, N. Photocatalysis in Chemical Biology: Extending the Scope of Optochemical Control and Towards New Frontiers in Semisynthetic Bioconjugates and Biocatalysis. *Helv. Chim. Acta* **2021**, *104*, e202100179.

(50) Biegasiewicz, K. F.; Cooper, S. J.; Gao, X.; Oblinsky, D. G.; Kim, J. H.; Garfinkle, S. E.; Joyce, L. A.; Sandoval, B. A.; Scholes, G. D.; Hyster, T. K. Photoexcitation of Flavoenzymes Enables a Stereoselective Radical Cyclization. *Science.* **2019**, *364*, 1166–1169.

(51) Crisenza, G. E. M.; Melchiorre, P. Chemistry Glows Green with Photoredox Catalysis. *Nat. Commun.* **2020**, *11*, 803.

(52) Sun, Z.; Fridrich, B.; De Santi, A.; Elangovan, S.; Barta, K. Bright Side of Lignin Depolymerization: Toward New Platform Chemicals. *Chem. Rev.* **2018**, *118*, 614–678.

(53) Li, S. H.; Liu, S.; Colmenares, J. C.; Xu, Y. J. A Sustainable Approach for Lignin Valorization by Heterogeneous Photocatalysis. *Green Chem.* **2016**, *18*, 594–607.

(54) Nguyen, S. T.; Murray, P. R. D.; Knowles, R. R. Light-Driven Depolymerization of Native Lignin Enabled by Proton-Coupled Electron Transfer. *ACS Catal.* **2020**, *10*, 800–805.

(55) Yayla, H. G.; Wang, H.; Tarantino, K. T.; Orbe, H. S.; Knowles, R. R. Catalytic Ring-Opening of Cyclic Alcohols Enabled by PCET Activation of Strong O-H Bonds. *J. Am. Chem. Soc.* **2016**, *138*, 10794–10797.

(56) Zhao, K.; Yamashita, K.; Carpenter, J. E.; Sherwood, T. C.; Ewing, W. R.; Cheng, P. T. W.; Knowles, R. R. Catalytic Ring Expansions of Cyclic Alcohols Enabled by Proton-Coupled Electron Transfer. J. Am. Chem. Soc. **2019**, 141, 8752–8757.

(57) Bosque, I.; Magallanes, G.; Rigoulet, M.; Kärkäs, M. D.; Stephenson, C. R. J. Redox Catalysis Facilitates Lignin Depolymerization. *ACS Cent. Sci.* **2017**, *3*, 621–628.

(58) Geyer, R.; Jambeck, J. R.; Law, K. L. Production, Use, and Fate of All Plastics Ever Made. *Sci. Adv.* **2017**, *3*, 25–29.