

Aryl amination using ligand-free Ni(II) salts and photoredox catalysis

Emily B. Corcoran,¹ Michael T. Pirnot,² Shishi Lin,³ Spencer D. Dreher,³ Daniel A. DiRocco,³ Ian W. Davies,³ Stephen L. Buchwald,^{2*} David W. C. MacMillan^{1*}

¹Merck Center for Catalysis at Princeton University, Princeton, NJ 08544, USA. ²Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA. ³Department of Process Chemistry, Merck Research Laboratories, Rahway, NJ 07065, USA.

*Corresponding author. E-mail: dmacmill@princeton.edu (D.W.C.M); sbuchwal@mit.edu (S.L.B.)

Over the past two decades, there have been major developments in transition metal-catalyzed aminations of aryl halides to form anilines, a common structure found in drug agents, natural product isolates, and fine chemicals. Many of these approaches have enabled highly efficient and selective coupling through the design of specialized ligands, which facilitate reductive elimination from a destabilized metal center. We postulated that a general and complementary method for C–N bond formation could be developed through the destabilization of a metal amido complex via photoredox catalysis, thus providing an alternative approach to the use of structurally complex ligand systems. Herein, we report the development of a distinct mechanistic paradigm for aryl amination using ligand-free nickel(II) salts, in which facile reductive elimination from the nickel metal center is induced via a photoredox-catalyzed electron-transfer event.

Transition metal-catalyzed cross-coupling reactions are fundamental methods for constructing complex molecular architectures. Key among these reactions has been the formation of carbon–nitrogen bonds to access anilines, a common structure found in medicinal agents and other complex molecules, through the coupling of amines with aryl halides or pseudohalides (1–6). Over the past two decades, multiple generations of ligands have been designed that have elevated palladium-catalyzed aryl amination to an essential transformation with both predictable reactivity and practical significance. The ligand frameworks that have been developed for palladium catalysis often employ a few key design elements (i.e., sterically encumbered architecture, hemilabile coordinating groups, electronically-tuned substituents) to enact high catalytic efficiency (7). One essential feature in ligand design is the capacity for destabilization of the palladium(II) amido complex, which induces reductive elimination resulting in C–N bond formation (8–10).

The broad success of ligated-palladium catalysis has hinged on the ability to selectively design, synthesize, and implement ligand classes to enable specific coupling reactions in high levels of efficiency and selectivity. Inspired by the effectiveness of structurally diverse ligand systems on palladium reductive elimination, we recognized that a general and complementary approach for productive C–N bond formation could be developed through an alternative mechanistic approach. In particular, we envisioned that the destabilization of a metal amido species could occur through

an electron transfer pathway via photoredox catalysis, thus inducing facile carbon–nitrogen bond formation analogous to the role of ligand in traditional palladium couplings (Fig. 1).

We proposed that nickel could be an ideal transition metal for this catalytic platform due to its known proclivity to participate in one-electron processes (11–13). Furthermore, this ligand-free mechanism could potentially broaden the applicability of nickel catalysis in aryl amination, (14–18) a system that is often plagued with the use of harsh reductants and air-sensitive nickel(0) complexes. Nickel(II) amido complexes, as detailed by Hillhouse and others, tend not to undergo reductive elimination at ambient temperature (19–21). Computations suggest that C–N reductive elimination from nickel(II) alkyl amido complexes is thermodynamically disfavored, in contrast to the exothermic reductive elimination from analogous palladium(II) alkyl amido complexes (22). Exposure of nickel(II) amido complexes to various oxidants results in facile reductive elimination, presumably through single-electron oxidation of the nickel(II) amido complex to the respective nickel(III) complex, lending support to our design strategy (19, 20).

In recent years, the merging of photoredox catalysis with nickel catalysis has enabled both C–O (23) and C–S (24, 25) bond formation through the generation of reactive radical intermediates and/or the single-electron modulation of nickel oxidation states (26, 27). The feasibility of these reactions arises from the capacity of photocatalysts to act as both strong single-electron oxidants and reductants upon

irradiation with visible light (28). Critical to the success of our design plan, the photoredox catalyst would be involved in two key aspects of our postulated mechanism: i) reduction of a nickel(II) salt to the active Ni(0) catalyst through the use of a mild sacrificial reductant and ii) electron transfer to destabilize the nickel(II) amido complex toward reductive elimination in the absence of an exogenous ligand.

In our proposed mechanism (Fig. 2), irradiation of heteroleptic iridium(III) photocatalyst $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$ [$\text{dF}(\text{CF}_3)\text{ppy}$ = 2-(2,4-difluorophenyl)-5-(trifluoromethyl)pyridine; dtbbpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine] (**1**) with visible light would produce long-lived triplet photoexcited state $^*\text{Ir}^{\text{III}}$ **2** (τ = 2.3 μs) (29). Concomitantly, nickel(0) complex **3** would oxidatively insert into bromoarene **4** to yield the corresponding nickel(II) aryl bromide complex **5**. The nickel(II) complex would then undergo ligand exchange and subsequent deprotonation to arrive at Ni(II)-aryl amido complex **6**. This nickel(II) complex could then participate in a single-electron transfer (SET) event with the photoexcited $^*\text{Ir}^{\text{III}}$ catalyst ($E_{1/2}^{\text{red}} [^*\text{Ir}^{\text{III}}/\text{Ir}^{\text{II}}] = +1.21$ V versus saturated calomel electrode (SCE) in CH_3CN) to afford Ni(III) complex **7** and the reduced iridium(II) photocatalyst (**8**) (29). Reductive elimination of the resultant nickel(III) complex would yield nickel(I) bromide species **9** and desired aniline product **10**. Both the photoredox and nickel catalytic cycles are closed upon a SET event between Ni(I)Br species **9** and Ir(II) ($E_{1/2}^{\text{red}} [\text{Ir}^{\text{III}}/\text{Ir}^{\text{II}}] = -1.37$ V versus SCE in CH_3CN) in which nickel(0) catalyst **3** and ground state iridium(III) catalyst **1** are regenerated simultaneously (29). The homogeneous nickel species is presumably coordinated to solvent, base, and/or amine throughout the process (denoted as L_n in Fig. 2), but would not require a designed ligand for efficient reactivity.

With this mechanistic hypothesis in hand, we first examined the proposed cross-coupling reaction with 4-bromobenzotrifluoride, pyrrolidine, nickel(II) bromide glyme, and a variety of ligands, bases, and solvents (see Supplementary Materials for details). The employment of photocatalyst **1**, 1,4-diazobicyclo[2.2.2]nonane (DABCO) as a base, and di-*tert*-butyl-2,2'-bipyridine as a supporting ligand provided the respective aniline product **10** in 72% yield in the presence of blue light. Control experiments established the importance of both nickel and photocatalyst, as no formation of the desired cross-coupled product was observed in the absence of either nickel, photocatalyst, or light. We sought to investigate the effectiveness of this protocol in the absence of ligand to confirm our initial mechanistic hypothesis. The cross-coupling reaction proceeded with enhanced reactivity in the absence of di-*tert*-butyl-2,2'-bipyridine, delivering the desired aniline product in 96% yield. This result supports our proposed mechanistic path-

way in Fig. 2 and the capacity of visible light and a photocatalyst to initiate reductive elimination in the absence of ligand.

With the optimal conditions in hand, we sought to demonstrate the generality of this ligand-free amination. As highlighted in Fig. 3, a wide variety of functional groups were tolerated on the amine coupling partner, including protected nitrogen atoms, trifluoromethyl groups, alcohols, alkenes, and sulfonamides (**11–15**, 72–77% yield). The high efficiency of the photoredox catalytic cycle is evidenced by the low photocatalyst loadings (≤ 0.02 mol%) required to effect the desired cross-coupling reactions. Monofluorinated amines (**16**) coupled cleanly under these conditions in 81% yield. Coupling of amine hydrogen chloride salts was accomplished in good yield by utilizing MTBD (7-methyl-1,5,7-diazabicyclo-[4.4.0]deca-5-ene) as the base (**16** and **17**, 81% and 70% yield); although the expense of MTBD often limits its practicality, synthetically useful yields were also obtained with DABCO in most cases (see Supplementary Materials for details). Chemoselective *N*-arylation in the presence of alcohols was also achieved, as no C–O coupling was observed (**13**, 76% yield, single regioisomer).

We found that α -substitution on the amine partner was tolerated in moderate yields (**20** and **22**, 60% and 78% yield, respectively), but hindered amines, *e.g.*, *tert*-butylamine, did not productively couple. Although nucleophilic amines, such as pyrrolidine and morpholine, coupled in good efficiencies (**10** and **22**, 96% and 91% yield, respectively) at ambient temperature after only a few hours, less nucleophilic substrates, such as trifluoroethylamine (**12**, 77% yield), allylamine (**14**, 76% yield), and furfurylamine (**23**, 90% yield), required longer reaction times (>24 hours). Amines lacking α -hydrogens failed to react under these reaction conditions. We hypothesized that an initial β -hydride elimination event is prerequisite for accessing the active Ni(0) catalyst in this reaction. Indeed, the addition of a substoichiometric amount of amine possessing α -hydrogens, such as pyrrolidine, to the reaction resulted in good yields (72% and 84% yield) of cross-coupled products **15** and **24**, respectively, with less than 10% of the pyrrolidine-coupled product being observed. For a number of substrates, product formation was observed in high yield utilizing $\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$ as the photocatalyst (see fig. S8). In cases where the yield is lower, protodehalogenation, along with trace phenol, aryl ether, or aryl chloride formation, can account for the mass balance of the reaction.

A variety of aryl bromides containing diverse functional groups, such as nitriles, amides, trifluoromethyl groups, halides, esters, and sulfonamides, performed well under the reaction conditions (**25–31**, 70–93% yield), with substitution at the *ortho* position being tolerated (**26** and **32**, 83% and 73% yield, respectively) (Fig. 3). Six-membered heteroar-

matic substrates were effectively coupled in good yields (**27** and **33–35**, 71–91% yield). Unfortunately, a series of aryl bromides, e.g. select heterocycles and a 2,6-disubstituted arene, failed to couple under the current reaction conditions (see fig. S8). For electron-rich arenes (**32** and **36**), lowering the photocatalyst loading (0.002 mol%) increased reaction efficiency and inhibited competitive protodehalogenation, as did the utilization of MTBD as base. Additionally, the initial evaluation of this C–N cross-coupling platform in flow chemistry has shown high efficiency in decreased reaction times (see Supplementary Materials).

To investigate the role of the photocatalyst in switching on nickel-catalyzed aryl amination, Ni(cod)₂ was employed. In the absence of light, the formation of aniline **10** was severely diminished (10% yield), while standard reaction conditions yielded **10** in 91% yield. This result supports our hypothesis that the cross-coupling is light-mediated and the effect of the photoredox catalyst is not limited to the reduction of a Ni(II) salt to the active Ni(0) species.

Lastly, this C–N cross-coupling platform was evaluated in a high throughput fashion using a microscale informer plate as developed by Merck laboratories. A chemistry informer library of eighteen complex, drug-like aryl halides was subjected to Ni-photoredox based amination conditions with piperidine. Remarkably, 78% of substrates underwent successful cross-coupling and provided “reaction hits” that were suitable for optimization (Fig. 4) (**30**). Benchmarked against other protocols that have been evaluated with this informer series, this study represents one of the most successful (**31**), generic catalysis platforms for aryl amination that has been evaluated within Merck. Based on these results, we are confident that this complementary method will find broad applicability in the synthesis of structurally diverse, drug-like compounds.

Through the merger of photoredox and nickel catalysis, challenging C–N reductive elimination can be switched on through the use of visible light and a photocatalyst via the intermediacy of a Ni(III) oxidation state (**32–34**). This strategy represents a complementary approach to traditional ligated-palladium catalysis through the use of a distinct mechanistic pathway for reductive elimination, which will likely be broadly applicable across a range of substrate classes.

REFERENCES AND NOTES

1. F. Ullmann, Ueber eine neue Bildungsweise von Diphenylaminderivaten. *Chem. Ber.* **36**, 2382–2384 (1903). doi:10.1002/cber.190303602174
2. I. Goldberg, Ueber Phenylirungen bei Gegenwart von Kupfer als Katalysator. *Chem. Ber.* **39**, 1691–1692 (1906). doi:10.1002/cber.19060390298
3. M. Kosugi, M. Kameyama, T. Migita, Palladium-catalyzed aromatic amination of aryl bromides with N,N-di-ethylamino-tributyltin. *Chem. Lett.* **12**, 927–928 (1983). doi:10.1246/cl.1983.927
4. M. S. Driver, J. F. Hartwig, A second-generation catalyst for aryl halide amination: Mixed secondary amines from aryl halides and primary amines catalyzed by (DPPF)PdCl₂. *J. Am. Chem. Soc.* **118**, 7217–7218 (1996). doi:10.1021/ja960937t
5. D. Chan, Promotion of reaction of N–H boths with triarylbiismuth and cupric acetate. *Tetrahedron Lett.* **37**, 9013–9016 (1996). doi:10.1016/S0040-4039(96)02116-8
6. A. S. Guram, S. L. Buchwald, Palladium-catalyzed aromatic aminations with in situ generated aminostannanes. *J. Am. Chem. Soc.* **116**, 7901–7902 (1994). doi:10.1021/ja00096a059
7. D. S. Surry, S. L. Buchwald, Dialkylbiaryl phosphines in Pd-catalyzed amination: A user’s guide. *Chem. Sci.* **2**, 27–50 (2011). doi:10.1039/C0SC00331J Medline
8. B. P. Fors, D. A. Watson, M. R. Biscoe, S. L. Buchwald, A highly active catalyst for Pd-catalyzed amination reactions: Cross-coupling reactions using aryl mesylates and the highly selective monoarylation of primary amines using aryl chlorides. *J. Am. Chem. Soc.* **130**, 13552–13554 (2008). doi:10.1021/ja8055358 Medline
9. J. F. Hartwig, Palladium-catalyzed amination of aryl halides: Mechanism and rational catalyst design. *Synlett* **1997**, 329–340 (1997). doi:10.1055/s-1997-789
10. A. T. Brusoe, J. F. Hartwig, Palladium-catalyzed arylation of fluoroalkylamines. *J. Am. Chem. Soc.* **137**, 8460–8468 (2015). doi:10.1021/jacs.5b02512 Medline
11. T. T. Tsou, J. K. Kochi, Mechanism of oxidative addition. Reaction of nickel(0) complexes with aromatic halides. *J. Am. Chem. Soc.* **101**, 6319–6332 (1979). doi:10.1021/ja00515a028
12. There is a report regarding the use of ligated nickel and photoredox catalysis for C–N bond formation (13).
13. S. Z. Tasker, T. F. Jamison, Highly regioselective indoline synthesis under nickel/photoredox dual catalysis. *J. Am. Chem. Soc.* **137**, 9531–9534 (2015). doi:10.1021/jacs.5b05597 Medline
14. J. P. Wolfe, S. L. Buchwald, Nickel-catalyzed amination of aryl chlorides. *J. Am. Chem. Soc.* **119**, 6054–6058 (1997). doi:10.1021/ja964391m
15. C.-Y. Gao, L.-M. Yang, Nickel-catalyzed amination of aryl tosylates. *J. Org. Chem.* **73**, 1624–1627 (2008). doi:10.1021/jo7022558 Medline
16. N. H. Park, G. Teverovskiy, S. L. Buchwald, Development of an air-stable nickel precatalyst for the amination of aryl chlorides, sulfamates, mesylates, and triflates. *Org. Lett.* **16**, 220–223 (2014). doi:10.1021/ol403209k Medline
17. S. Ge, R. A. Green, J. F. Hartwig, Controlling first-row catalysts: Amination of aryl and heteroaryl chlorides and bromides with primary aliphatic amines catalyzed by a BINAP-ligated single-component Ni(0) complex. *J. Am. Chem. Soc.* **136**, 1617–1627 (2014). doi:10.1021/ja411911s Medline
18. T. Shimasahi, M. Tobisu, N. Chatani, Nickel-catalyzed amination of aryl pivalates by the cleavage of aryl C–O bonds. *Angew. Chem. Int. Ed.* **49**, 2929 (2010). doi:10.1002/anie.200907287
19. K. Koo, G. L. Hillhouse, Carbon-nitrogen bond formation by reductive elimination from nickel(II) amido alkyl complexes. *Organometallics* **14**, 4421–4423 (1995). doi:10.1021/om00009a054
20. B. L. Lin, C. R. Clough, G. L. Hillhouse, Interactions of aziridines with nickel complexes: Oxidative-addition and reductive-elimination reactions that break and make C–N bonds. *J. Am. Chem. Soc.* **124**, 2890–2891 (2002). doi:10.1021/ja017652n Medline
21. L. Lies, T. Matsubara, E. Nakamura, Nickel-catalyzed synthesis of diarylamines via oxidatively induced C–N bond formation at room temperature. *Org. Lett.* **14**, 5570–5573 (2012). doi:10.1021/ol302688u Medline
22. S. A. Macgregor, G. W. Neave, C. Smith, Theoretical studies on C-heteroatom bond formation via reductive elimination from group 10 M(PH₃)₂(CH₃)(X) species (X = CH₃, NH₂, OH, SH) and the determination of metal–X bond strengths using density functional theory. *Faraday Discuss.* **124**, 111–127, discussion 145–153, 453–455 (2003). doi:10.1039/B212309F Medline
23. J. A. Terrett, J. D. Cuthbertson, V. W. Shurtleff, D. W. C. MacMillan, Switching on elusive organometallic mechanisms with photoredox catalysis. *Nature* **524**, 330–334 (2015). doi:10.1038/nature14875 Medline
24. M. S. Oderinde, M. Frenette, D. W. Robbins, B. Aquila, J. W. Johannes, Photoredox mediated nickel catalyzed cross-coupling of thiols with aryl and heteroaryl iodides via thiyl radicals. *J. Am. Chem. Soc.* **138**, 1760–1763 (2016). doi:10.1021/jacs.5b11244 Medline
25. M. Jouffroy, C. B. Kelly, G. A. Molander, Thioetherification via photoredox/nickel dual catalysis. *Org. Lett.* **18**, 876–879 (2016). doi:10.1021/acs.orglett.6b00208 Medline
26. J. C. Tellis, D. N. Primer, G. A. Molander, Dual catalysis. Single-electron transmetalation in organoboron cross-coupling by photoredox/nickel dual